arXiv:1002.4285v2 [math-ph] 13 Jul 2010

Complex and biHermitian structures on four dimensional real Lie algebras

A. Rezaei-Aghdam* and M. Sephid†

Department of Physics, Faculty of science, Azarbaijan University
of Tarbiat Moallem , 53714-161, Tabriz, Iran

July 14, 2010

Abstract

We give a new method for calculation of complex and biHermitian structures on low dimensional real Lie algebras. In this method, using non-coordinate basis, we first transform the Nijenhuis tensor field and biHermitian structure relations on Lie groups to the tensor relations on their Lie algebras. Then we use adjoint representation for writing these relations in the matrix form; in this manner by solving these matrix relations and using automorphism groups of four dimensional real Lie algebras we obtain and classify all complex and biHermitian structures on four dimensional real Lie algebras.

---

*e-mail: rezaei-a@azaruniv.edu

†e-mail: s.sephid@azaruniv.edu

1 Introduction

Calculation of complex structures on homogeneous complex manifolds and especially on Lie groups is important from both mathematical and physical point of view. Mathematically, classification of these manifolds is based on determination of possible complex structures. From the physical point of view these structures have an important role in the N=(2,2) supersymmetric sigma models [1]. It is shown that the N=(2,2) extended supersymmetry in sigma model implies the existence of biHermitian structure on the target manifold such that the complex structures are covariantly constant with respect to torsionful affine connections (see for example [2] and references therein). Furthermore, it is shown that the algebraic structures related to these biHermitian structures for N=(2,2) supersymmetric WZW models are the Manin triples [3],[4]. For these reasons, the calculation of the complex and the biHermitian structures on manifolds especially on Lie groups, are important. Samelson [5] shows that compact Lie groups always admit an invariant complex structure . As for the non-compact case, Morimoto [6] proves that there always exist invariant complex structures on any even dimensional reductive Lie groups. In [7] and [8], complex structures on real four dimensional Lie algebras are classified. The method used in this works is special and seems not to be adequate for the calculation in higher dimensions. In the present paper we give a new method for this purpose, which can be applied for low dimensional Lie groups . In this method, using non-coordinate basis, we first transform the Nijenhuis tensor on Lie groups to algebraic tensor relations on their Lie algebras. Then, using adjoint representation we rewrite these relations in the matrix form. Finally, we solve these matrix relations using Maple. In this research we perform this for real four dimensional Lie algebras. The results for some algebras are different and complete with respect to [8]. Furthermore, calculation of biHermitian structure for four dimensional Lie algebras is new .

The paper is organized as follows. In section 2, using non-coordinate basis, we transform the Nijenhuis tensor relation on a Lie group to the algebraic tensor relation on its Lie algebra. Then, using adjoint representation, we write these relations in the matrix form. The relations can also be obtained from definition of complex structures on Lie algebras as endomorphism of them. Then, in section 3 using Maple we solve these matrix relations to obtain complex structures on real four dimensional Lie algebras. In this process, we apply automorphism groups of real four dimensional Lie algebras for obtaining non-equivalent complex structures (table 1). We then compare our results with [7] and [8]. Note that here we use Patera *etal* [9] classification of real four dimensional Lie algebras. The list of Lie algebras and their automorphism groups [10] are given in appendix. In section 4, we first transform the tensorial form of the biHermitian relations on Lie groups into the algebraic tensorial relations on their Lie algebras. In this respect, we define biHermitian structure on Lie algebra independently and give an equivalent relation for obtaining non equivalent biHermitian structures. Then using adjoint representation we rewrite these relations in the matrix forms and solve them by Maple. Therefore, the present paper will be a continuation to the discussion of biHermitian structure on real four dimensional Lie algebras. Some discussions are given in the conclusion section.

2 A brief review of complex structures on Lie groups

Definition 1: *Let M be a differentiable manifold, then the pair (M,J) is called almost complex manifold if there exists a tensor field J of type (1,1) such that at each point p of M ,* $J_p^2 = -1$*; tensor field J is also called the almost complex structure. Furthermore, if the Lie bracket of any vector fields of type* $(1,0)$ $X, Y \in T_pM^+$ *is again of the same type, then the complex structure* J_p *is said to be integrable, where* $T_pM^+ = \{Z \in T_pM^C \mid J_pZ = +iZ\}$.

Theorem (Newlander and Nirenberg [11]) : *An almost complex structure J on a manifold M is integrable if and only if*

$$N(X,Y) = 0 \qquad \forall X, Y \in \chi(M) \,, \tag{1}$$

where $\chi(M)$ *is the set of vector fields on M and the Nijenhuis tensor* $N : \chi(M) \otimes \chi(M) \to \chi(M)$ *is given by*

$$N(X,Y) = [X,Y] + J[JX,Y] + J[X,JY] - [JX,JY] \,. \tag{2}$$

In the coordinate basis, i.e the basis $\{e_\mu = \frac{\partial}{\partial x^\mu}\}$ and $\{dx^\mu\}$ for vectors and dual vectors (forms) respectively on M, the almost complex structures and Nijenhuis tensor are expressed as $J = J_\mu{}^\nu e_\nu \otimes dx^\mu$ and $N = N_{\mu\nu}{}^\lambda dx^\mu \otimes dx^\nu \otimes e_\lambda$ respectively and the integrability condition (2) can be rewritten as follows :

$$N_{\kappa\nu}{}^\mu = J_\lambda{}^\mu(\partial_\kappa J_\nu{}^\lambda) + J_\nu{}^\lambda(\partial_\lambda J_\kappa{}^\mu) - J_\lambda{}^\mu(\partial_\nu J_\kappa{}^\lambda) - J_\kappa{}^\lambda(\partial_\lambda J_\nu{}^\mu \;= 0. \tag{3}$$

Meanwhile, the relation $J^2 = -1$ can be rewritten as

$$J_{\lambda}{}^{\mu} J_{\mu}{}^{\nu} = -\delta_{\lambda}{}^{\nu}. \tag{4}$$

Furthermore one can rewrite the above equations using the non-coordinate basis $\{\hat{e}_\alpha\}$ and $\{\hat{\theta}^\alpha\}$ for vectors and forms on M. For these basis we have

$$\hat{e}_\alpha = e_{\alpha}{}^{\mu} e_\mu \qquad , \qquad e_{\alpha}{}^{\mu} \in GL(m, R) \ , \tag{5}$$

where for the vierbeins $e_{\alpha}{}^{\mu}$ and its inverse $e^{\alpha}{}_{\mu}$ we have

$$e^{\alpha}{}_{\mu} e_{\beta}{}^{\mu} = \delta^{\alpha}{}_{\beta} \ \ , \ \ e^{\alpha}{}_{\mu} e_{\alpha}{}^{\nu} = \delta_{\mu}{}^{\nu} \ . \tag{6}$$

The dual basis $\{\hat{\theta^\alpha}\}$ are defined by $< \hat{\theta^\alpha}, \hat{e}_\beta >= \delta^{\alpha}{}_{\beta}$ and we have $\hat{\theta^\alpha} = e^{\alpha}{}_{\mu} dx^\mu$. Furthermore, the vierbeins satisfy the following relation:

$$f_{\alpha\beta}{}^{\gamma} = e^{\gamma}{}_{\nu}(e_{\alpha}{}^{\mu}\partial_\mu e_{\beta}{}^{\nu} - e_{\beta}{}^{\mu}\partial_\mu e_{\alpha}{}^{\nu}) \ , \tag{7}$$

if M is a Lie group manifold G then $f_{\alpha\beta}{}^{\gamma}$,s are structure constants of Lie algebra **g** of G. Now on these bases the tensor $J = J_{\mu}{}^{\nu} e_\nu \otimes dx^\mu$ can be rewritten as

$$J_{\mu}{}^{\nu} = e^{\alpha}{}_{\mu} J_{\alpha}{}^{\beta} e_{\beta}{}^{\nu} \ , \tag{8}$$

where $J_{\alpha}{}^{\beta}$ is an endomorphism of **g**, i.e. $J : \mathbf{g} \longrightarrow \mathbf{g}$. Now by applying this relation to (4) we have the following matrix relation for matrices $J_{\alpha}{}^{\beta}$:

$$J^2 = -I \ . \tag{9}$$

Furthermore, by applying relations (7) and (8) to tensor equation (3) and using (6) and assuming that $J_{\alpha}{}^{\beta}$ and $g_{\alpha\beta}$ are independent of coordinates of G, after some calculations we have the following algebraic relation for(3):

$$f_{\beta\alpha}{}^{\gamma} + J_{\sigma}{}^{\gamma}\ J_{\alpha}{}^{\delta}\ f_{\beta\delta}{}^{\sigma} - J_{\beta}{}^{\sigma}\ J_{\alpha}{}^{\delta}\ f_{\sigma\delta}{}^{\gamma} + J_{\beta}{}^{\sigma}\ J_{\delta}{}^{\gamma}\ f_{\sigma\alpha}{}^{\delta} = 0. \tag{10}$$

Finally, using adjoint representations

$$f_{\beta\alpha}{}^{\gamma} = -(\mathcal{Y}^{\gamma})_{\beta\alpha} \quad , \quad f_{\beta\alpha}{}^{\gamma} = -(\chi_\beta)_{\alpha}{}^{\gamma}, \tag{11}$$

the relation (10) will have the following matrix form:

$$\mathcal{Y}^{\alpha} + J\ \mathcal{Y}^{\beta}\ J_{\beta}{}^{\alpha} + J_{\beta}{}^{\alpha}\ \mathcal{Y}^{\beta}\ J^t - J\ \mathcal{Y}^{\alpha}\ J^t = 0 \ , \tag{12}$$

or

$$\chi_\alpha + J\ \chi_\alpha\ J + J_{\alpha}{}^{\beta}\ \chi_\beta\ J - J_{\alpha}{}^{\beta}\ J\ \chi_\beta = 0 \ . \tag{13}$$

Note that the above equation can also be obtained from the definition of complex structure on Lie algebra **g** as follows.

Definition 2: *An integrable complex structure on a real Lie algebra* **g** *is an endomorphism J of* **g** *such that*

a) $J^2 = -Id,$ (14)

b) $[X, Y] + J[JX, Y] + J[X, JY] - [JX, JY] = 0 \ , \qquad \forall X, Y \in \mathbf{g}$. (15)

Now if we use $\{X_\alpha\}$ as basis for Lie algebra **g** with the following structure constants:

$$[X_\alpha, X_\beta] = f_{\alpha\beta}{}^{\gamma} X_\gamma \ , \tag{16}$$

and use the following relation for J:

$$J X_\alpha = J_{\alpha}{}^{\beta} X_\beta \ , \tag{17}$$

then relations (14) and (15) can be rewritten as (9), (12) or (13). Now, in order to obtain algebraic complex structures $J_{\alpha}{}^{\beta}$, it is enough to solve equations (9) and (12) or (13) simultaneously . We do this for real four dimensional Lie algebras in the next section.

3 Calculation of complex structures on four dimensional real Lie algebras

In this section we use the Patera *etal* classification [9] of four dimensional real Lie algebras . The commutation relations and the automorphism groups of these Lie algebras are given in the appendix [1]. Now one can write the adjoint representation ($\mathcal{Y}$) for these Lie algebras and then solve the matrix relations (9) and (12) for obtaining complex structures. We do this by Maple. Note that in this process one can obtain equivalent complex structures; to avoid these and in order to obtain inequivalent complex structure we use the following equivalent relation:

Definition 3[**8**] *: Two complex structures J_1 and J_2 of Lie algebra* $\mathbf{g}$ *are equivalent if there exists an element A of automorphism group of Lie algebra* $\mathbf{g}$ *($Aut(\mathbf{g})$) such that:*

$$J_2 = A \; J_1 \; A^{-1} \; . \tag{18}$$

Note that this relation is an equivalent relation.
In this way, we do this and obtain all non-equivalent complex structures on four dimensional real Lie algebras. The results are classified in table 1 [2]. As the table shows 21 out of 30 real four dimensional Lie algebras have complex structures. To compare these results with results of [8] first we must obtain the isomorphism relations between the four dimensional real Lie algebras presented in [9] and those presented in [8]. According to the calculations in [8] and [12] we have isomorphism relations as summarized in the following table:

$4A_1$	$A_{4,1}$	$A^a_{4,2}$	$A_{4,3}$	$A_{4,4}$	$A^{a,b}_{4,5}$	$A^{a,b}_{4,6}$	$A_{4,7}$	$A_{4,8}$	$A^b_{4,9}$	$A_{4,10}$	$A^a_{4,11}$	$A_{4,12}$
$\mathfrak{a}_4$	$\mathfrak{n}_4$	$\mathfrak{r}_{4,a}$	$\mathfrak{r}_{4,0}$	$\mathfrak{r}_4$	$\mathfrak{r}_{4,a,b}$	$\mathfrak{r}'_{4,a,b}$	$\mathfrak{h}_4$	$\mathfrak{d}_4$	$\mathfrak{d}_{4,1/1+b}$	$\mathfrak{d}'_{4,0}$	$\mathfrak{d}'_{4,a}$	$\mathfrak{aff}(\mathbb{C})$

$A_2 \oplus A_2$	$II \oplus R$	$III \oplus R$	$IV \oplus R$	$V \oplus R$	$VI_0 \oplus R$	$VI_a \oplus R$ $(a \neq 0,1)$	$VII_0 \oplus R$	$VII_a \oplus R$ $(a \neq 0)$
$\mathfrak{r}_2\mathfrak{r}_2$	$\mathfrak{r}\mathfrak{h}_3$	$\mathfrak{r}\mathfrak{r}_{3,0}$	$\mathfrak{r}\mathfrak{r}_3$	$\mathfrak{r}\mathfrak{r}_{3,1}$	$\mathfrak{r}\mathfrak{r}_{3,-1}$	$\mathfrak{r}\mathfrak{r}_{3,a}$	$\mathfrak{r}\mathfrak{r}'_{3,0}$	$\mathfrak{r}\mathfrak{r}'_{3,a}$

In this respect, one can see that in [8] one complex structure is obtained for Lie algebra $VII_0 + R$ but according to our calculation this Lie algebra has two non-equivalent complex structures. For non-solvable Lie algebras $VIII + R$ and $IX + R$ we obtain complex structures. In [8], two complex structures are obtained for Lie algebra $A_{4,8}$ but here we obtain four complex structures to this Lie algebra. Furthermore, two complex structures are obtained for Lie algebra $A_{4,10}$ in [8], but according to table 1 we obtain four complex structures for this Lie algebra. Meanwhile, in [8], two complex structures are obtained for Lie algebra $A_{4,12}$ but here we obtain three complex structures for this Lie algebra. The results for other Lie algebras are the same as [8].

---

[1] Note that for decomposable Lie algebra ($L_3 \oplus R$) we use Bianchi classification of real three dimensional Lie algebras L_3.

[2] Note that in this table the bases are shown with $\{e_\alpha\}$ instead of $\{X_\alpha\}$

TABLE 1 : Complex structures on four dimensional real Lie algebras

Lie Algebra	Complex Structures
$III \oplus R$	$Je_1 = -e_2 \ , \ Je_3 = e_1 - e_4$
$A_2 \oplus A_2$	$Je_1 = -e_2 \ , \ Je_3 = -e_4$
$II \oplus R$	$Je_1 = -e_4 \ , \ Je_2 = e_3$
$V \oplus R$	$Je_1 = -e_4 \ , \ Je_2 = e_3$
$VII_0 \oplus R$	$J_1e_1 = -e_2 \ , \ J_1e_3 = -e_4$ $J_2e_1 = e_2 \ , \ J_2e_3 = -e_4$
$VII_a \oplus R$ $(a \neq 0)$	$J_1e_1 = -e_4 \ , \ J_1e_2 = e_3$ $J_2e_1 = -e_4 \ , \ J_2e_2 = -e_3$
$VIII \oplus R$	$Je_1 = -e_2 \ , \ Je_3 = pe_3 - (1+p^2)e_4 \qquad (p \in \mathbb{R})$
$IX \oplus R$	$Je_1 = -e_2 \ , \ Je_3 = pe_3 - (1+p^2)e_4 \qquad (p \in \mathbb{R})$
$A_{4,2}^1$	$Je_1 = e_2 \ , \ Je_3 = e_4$
$A_{4,5}^{a,a}$ $(-1 \leq a < 1, a \neq 0)$	$Je_1 = -e_4 \ , \ Je_2 = e_3$
$A_{4,5}^{a,1}$ $(-1 \leq a < 1, a \neq 0)$	$Je_1 = e_3 \ , \ Je_2 = e_4$
$A_{4,5}^{1,1}$	$Je_1 = -e_3 \ , \ Je_2 = -e_4$
$A_{4,6}^{a,b}$ $(a \neq 0, b \geq 0)$	$J_1e_1 = e_4 \ , \ J_1e_2 = e_3$ $J_2e_1 = e_4 \ , \ J_2e_2 = -e_3$
$A_{4,7}$	$Je_1 = e_2 \ , \ Je_3 = -e_4$
$A_{4,8}$	$J_1e_1 = e_2 \ , \ J_1e_3 = e_4$ $J_2e_1 = -e_3 \ , \ J_2e_2 = -e_4$ $J_3e_1 = e_2 \ , \ J_3e_3 = e_2 + e_4$ $J_4e_1 = -(e_1 + 2e_3) \ , \ J_4e_2 = -(e_1 + e_2) - 2(e_3 + e_4)$
$A_{4,9}^b$ $(0 <\mid b \mid< 1)$	$J_1e_1 = -be_2 \ , \ J_1e_3 = e_4$ $J_2e_1 = -e_3 \ , \ J_2e_2 = -e_4$
$A_{4,9}^1$	$J_1e_1 = -e_3 \ , \ J_1e_2 = -e_4$ $J_2e_1 = e_4 \ , \ J_2e_2 = -e_3$ $J_3e_1 = -e_4 \ , \ J_3e_2 = -e_3$
$A_{4,9}^0$	$Je_1 = -e_3 \ , \ Je_2 = -e_4$
$A_{4,10}$	$J_1e_2 = -e_1 + e_3 \ , \ J_1e_3 = -e_1 - e_2 + e_3 + e_4$ $J_2e_2 = e_1 + e_3 \ , \ J_2e_3 = -e_1 - e_2 - e_3 + e_4$ $J_3e_2 = -e_1 - e_3 \ , \ J_3e_3 = e_1 + e_2 + e_3 - e_4$ $J_4e_2 = -e_1 - e_3 \ , \ J_4e_3 = -e_1 + e_2 - e_3 + e_4$
$A_{4,11}^a$ $(a > 0)$	$J_1e_1 = e_4 \ , \ J_1e_2 = -e_3$ $J_2e_1 = e_4 \ , \ J_2e_2 = e_3$ $J_3e_1 = -e_4 \ , \ J_3e_2 = -e_3$ $J_4e_1 = -e_4 \ , \ J_4e_2 = e_3$
$A_{4,12}$	$J_1e_1 = e_2 \ , \ J_1e_3 = e_4$ $J_2e_1 = -e_2 \ , \ J_2e_3 = e_4$ $J_3e_1 = e_4 \ , \ J_3e_2 = e_3$

4 BiHermitian structures on four dimensional real Lie algebras

Definition 4 [1] [2] :*If the complex manifold M has two complex structures* $J_\pm$ *such that it is Hermitian with respect to both complex structures. i.e.*

$$J_\pm^2 = -1 \; , \tag{19}$$

$$N_{\mu\nu}{}^{\kappa}(J_\pm) = 0 \; , \tag{20}$$

$$J_{\pm\mu}{}^{\lambda} \; g_{\lambda\eta} \; J_{\pm\nu}{}^{\eta} = g_{\mu\nu}, \tag{21}$$

and furthermore if these complex structures be covariantly constant with respect to certain connections $\Gamma^\pm$

$$\bigtriangledown_\mu^\pm \quad J_{\pm\nu}{}^{\lambda} \equiv J_{\pm\nu}{}^{\lambda}{}_{,\mu} + \Gamma^{\pm}_{\mu\rho}{}^{\lambda} J_{\pm\nu}{}^{\rho} - \Gamma^{\pm}_{\mu\nu}{}^{\rho} J_{\pm\rho}{}^{\lambda} = 0, \tag{22}$$

with

$$\Gamma^{\pm}_{\mu\nu}{}^{\lambda} = \Gamma_{\mu\nu}{}^{\lambda} \pm T_{\mu\nu}{}^{\lambda} \;\; , \;\; T_{\mu\nu}{}^{\lambda} = H_{\mu\nu\eta} g^{\eta\lambda}, \tag{23}$$

then it is said that M has biHermitian structure, shown by $(M, g, J_\pm)$*.*

In the above definition $g_{\mu\nu}, \Gamma_{\mu\nu}{}^{\lambda}$ and $H_{\mu\nu\eta}$ are metric , Christoffel connection and antisymmetric tensors on M respectively. Using (22), the integrability condition (20) may be rewritten in the following form [2]:

$$H_{\delta\nu\lambda} = J_{\pm\delta}{}^{\sigma} J_{\pm\nu}{}^{\rho} H_{\sigma\rho\lambda} + J_{\pm\delta}{}^{\rho} J_{\pm\lambda}{}^{\sigma} H_{\sigma\rho\nu} + J_{\pm\upsilon}{}^{\sigma} J_{\pm\lambda}{}^{\rho} H_{\sigma\rho\delta} \; . \tag{24}$$

Furthermore, by introducing the Kähler forms

$$\omega_{\pm\mu\nu} \equiv \;\; g_{\mu\nu} J_{\pm\nu}{}^{\lambda}, \tag{25}$$

and by use of (22) one can find

$$(d\omega_\pm)_{\rho\mu\nu} = \pm(H_{\sigma\rho\mu} J^{\sigma}_{\pm\nu} + H_{\sigma\mu\nu} J^{\sigma}_{\pm\rho} + H_{\sigma\nu\rho} J^{\sigma}_{\pm\mu}), \tag{26}$$

where

$$(d\omega_\pm)_{\lambda\sigma\gamma} = \frac{1}{2}(\partial_\lambda \omega_{\pm\sigma\gamma} + \partial_\sigma \omega_{\pm\gamma\lambda} + \partial_\gamma \omega_{\pm\lambda\sigma}). \tag{27}$$

Finally, using (25) and (26) one can find

$$H_{\mu\nu\rho} = -J^{\lambda}_{+\mu} J^{\sigma}_{+\nu} J^{\gamma}_{+\rho} (d\omega_+)_{\lambda\sigma\gamma} = -J^{\lambda}_{-\mu} J^{\sigma}_{-\nu} J^{\gamma}_{-\rho} (d\omega_-)_{\lambda\sigma\gamma}. \tag{28}$$

In this respect, the target manifold $(M, g, J_\pm)$ is said to have biHermitian structure if two Hermitian complex structures $J_\pm$ satisfy the relation (28) (i.e. relation between (J_+, ω_+) and (J_-, ω_-))which defines the torsion H. Now, for the case where M is a Lie group G, similar to the process presented in section 2 one can transform relations $(19) - (22)$ and (24) to the algebraic relation using the relations $(6), (7)$ and the following relations:

$$g_{\alpha\beta} = L_\alpha{}^{\mu} L_\beta{}^{\nu} g_{\mu\nu} = R_\alpha{}^{\mu} R_\beta{}^{\nu} g_{\mu\nu} \;\; , \;\; g_{\mu\nu} = L^{\alpha}{}_{\mu} L^{\beta}{}_{\nu} g_{\alpha\beta} = R^{\alpha}{}_{\mu} R^{\beta}{}_{\nu} g_{\alpha\beta}, \tag{29}$$

$$H_{\mu\nu\rho} = \frac{1}{2} L^{\alpha}{}_{\mu} L^{\beta}{}_{\nu} L^{\gamma}{}_{\rho} H_{\alpha\beta\gamma} = \frac{1}{2} R^{\alpha}{}_{\mu} R^{\beta}{}_{\nu} R^{\gamma}{}_{\rho} H_{\alpha\beta\gamma}, \tag{30}$$

$$J_{+\mu}{}^{\nu} = R^{\alpha}{}_{\mu} J_\alpha{}^{\beta} R_\beta{}^{\nu} \; , \; J_{-\mu}{}^{\nu} = L^{\alpha}{}_{\mu} J_\alpha{}^{\beta} L_\beta{}^{\nu}, \tag{31}$$

where $L^{\alpha}{}_{\mu}(R^{\alpha}{}_{\mu})$ and $L_\beta{}^{\nu}(R_\beta{}^{\nu})$ are left(right) invariant vierbeins and their inverses respectively. Now using these relations , (21) and (24) transform to the following matrix relations:

$$J \; g \; J^t = g, \tag{32}$$

$$H_\alpha = J(H_\beta J_\alpha{}^{\beta}) + J H_\alpha J^t + (H_\beta J_\alpha{}^{\beta}) J^t, \tag{33}$$

where $(H_\alpha)_{\beta\gamma} = H_{\alpha\beta\gamma}$. Furthermore, using the following relations [13] [3]

$$\bigtriangledown^\rho L_\alpha{}^\mu = -\frac{1}{2}(f_\alpha{}^{(\rho\mu)} + f^{\rho\mu}{}_\alpha + T_\alpha{}^{(\rho\mu)} + T^{\rho\mu}{}_\alpha + L_\beta{}^\rho L_\gamma{}^\mu \bigtriangledown_\alpha g^{\beta\gamma} + L^{\beta\rho} \bigtriangledown^\mu g_{\alpha\beta} - L^{\beta\mu} \bigtriangledown^\rho g_{\alpha\beta}), \tag{34}$$

$$\bigtriangledown^\rho R_\alpha{}^\mu = -\frac{1}{2}(-f_\alpha{}^{(\rho\mu)} - f^{\rho\mu}{}_\alpha + T_\alpha{}^{(\rho\mu)} + T^{\rho\mu}{}_\alpha + R_\beta{}^\rho R_\gamma{}^\mu \bigtriangledown_\alpha g^{\beta\gamma} + R^{\beta\rho} \bigtriangledown^\mu g_{\alpha\beta} - R^{\beta\mu} \bigtriangledown^\rho g_{\alpha\beta}), \tag{35}$$

and assuming that $g_{\alpha\beta}$ are coordinate independent; the relation (22) transforms to the following algebraic relation[4]:

$$J(H_\alpha - \chi_\alpha g) = (J(H_\alpha - \chi_\alpha g))^t. \tag{36}$$

Note that the metric $g_{\alpha\beta}$ is the ad invariant metric on Lie algebras $\mathbf{g}$ .i.e. we have

$$\langle X_\alpha, X_\beta \rangle = g_{\alpha\beta}, \tag{37}$$

$$\langle X_\alpha, [X_\beta, X_\gamma] \rangle = \langle [X_\alpha, X_\beta], X_\gamma \rangle, \tag{38}$$

or in matrix notation we have [5]

$$\chi_\alpha g = -(\chi_\alpha g)^t. \tag{39}$$

Now, one can obtain biHermitian structures on Lie algebras by solving relations (9), (12), (32), (33), (36) and (39) simultaneously. These relations can be applied on the Lie algebra as a definition of algebraic biHermitian structure on $\mathbf{g}$;

Definition 5: *If there exist endomorphism $J : \mathbf{g} \rightarrow \mathbf{g}$ of Lie algebra with ad invariant metric g and antisymmetric bilinear map $H : \mathbf{g} \otimes \mathbf{g} \rightarrow \mathbf{g}$ such that the relations* (9), (12), (32), (33), (36) *and* (39) *are satisfied, then we have biHermitian structure (J, g, H) on* $\mathbf{g}$.

Note that relation (33) is equivalent to matrix relation of integrability condition i.e. relation (12) . For this reason, first it is better to obtain algebraic complex structures J, then solve relations (32), (36) and (39) and finally check them in (33). We do this for real four dimensional Lie algebras using Maple. Note that similar to complex structures, in order to obtain non-equivalent biHermitian structures we suggest the following equivalent relations.

Definition 6 *: Two biHermitian structures (J, g, H) and $(J^{'}, g^{'}, H^{'})$ of Lie algebras* $\mathbf{g}$ *are equivalent if there exists an element A of automorphism group of Lie algebra* $\mathbf{g}$ *(Auto* $\mathbf{g}$*) such that:*

$$J^{'} = AJA^{-1}, \tag{40}$$

$$g^{'} = AgA^t, \tag{41}$$

$$H^{'}_\alpha = A(H_\beta A_\alpha{}^\beta)A^t. \tag{42}$$

These relations are equivalent relations and are satisfied in the equivalent conditions. Note that if $f_{\beta\gamma}{}^\alpha = H_{\delta\beta\gamma} g^{\delta\alpha}$ or H is isomorphic with f, i.e. if there exists isomorphism matrix C such that

$$CY^\alpha C^t = \tilde{Y}^\beta C_\beta{}^\alpha, \tag{43}$$

where $(Y^\alpha)_{\beta\gamma} = -f_{\beta\gamma}{}^\alpha$ and $(\tilde{Y}^\alpha)_{\beta\gamma} = -H_{\delta\beta\gamma} g^{\delta\alpha}$; then (J, g, H) shows the Manin triple structure on $\mathbf{g}$[4]. In this way biHermitian structures on real four dimensional Lie algebras can be classified as table 2 . Note that according to the table for Lie algebra $A_{4,8}$, we have two non-equivalent biHermitian structure (J, g, H) where the second biHermitian structure shows the Manin triple structure of $A_{4,8}$ [4] (i.e. $A_{4,8}$ is a Manin triple of two dimensional Lie bialgebras (type B and semiabelian)[14]) for the following values of parameters:

---

[3] Note that in these relations all algebraic (target) indices are lowered and raised by $g_{\alpha\beta}(g_{\mu\nu})$; furthermore, these indices transform into each other by $L_\alpha{}^\mu(R_\alpha{}^\mu)$ or $L^\alpha{}_\mu(R^\alpha{}_\mu)$. The symmetrization notation have the following form: $f_\alpha{}^{(\rho\mu)} = f_\alpha{}^{\rho\mu} + f_\alpha{}^{\mu\rho}$.

[4] This relation can also be obtained from algebraic form of (28).

[5] For semisimple Lie algebras, the Killing form is one of nondegenerate solutions for this equation and other nondegenerate solution may exist. For nonsemisimple Lie algebras the Killing tensor degenerates and there may exist nondegenerate solution for (39).

$$c_1 = c_2 = c_3 = c_4 = c_5 = c_6 = c_7 = c_9 = c_{10} = c_{11} = c_{13} = c_{14} = 0 \ , \ c_{12} = c_{15} = -1.$$

For Lie algebras $VIII \oplus R$, there is one biHermitian structure where this structure, for the values

$$d_1 = d_2 = d_4 = d_5 = d_7 = d_8 = d_{10} = d_{11} = d_{12} = d_{14} = d_{15} = d_{16} = 0 \ , \ d_9 \neq 0 \ , \ \ d_3 = -d_6 = \alpha,$$

is isomorphic with two dimensional Lie bialgebra type A [14]. There exists one biHermitian structure for Lie algebra $IX \oplus R$. The results are given in table 2 [6]. Note that the isomorphism relation (43)(i.e. the biHermitian structures which show Manin triple) are independent of the choice of special biHermitian structures from equivalent class of biHermitian structures. In this way if relation (43) holds ; then by $\tilde{Y}'^{\alpha} = -H'_{\delta} g'^{\delta\alpha}$ and using relations (42) and (43) one can show that

$$(AC)Y^{\gamma}(AC)^t = \tilde{Y}'^{\alpha}(AC)_{\alpha}{}^{\gamma}. \tag{44}$$

TABLE 2 : biHermitian structures on four dimensional real Lie algebras

Lie Algebra	Complex Structures	g	antisymmetric tensor
$A_{4,8}$	$J = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}$	$g = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$	$H_1 = \begin{pmatrix} 0 & b_1 & 1 & -b_2 \\ -b_1 & 0 & b_2 & 1 \\ -1 & -b_2 & 0 & b_3 \\ b_2 & -1 & -b_3 & 0 \end{pmatrix}$ $H_2 = \begin{pmatrix} 0 & b_4 & b_5 & -b_6 \\ -b_4 & 0 & b_6 & b_5 \\ -b_5 & -b_6 & 0 & b_7 \\ b_6 & -b_5 & -b_7 & 0 \end{pmatrix}$ $H_3 = \begin{pmatrix} 0 & b_8 & b_9 & -b_{10} \\ -b_8 & 0 & b_{10} & 1+b_9 \\ -b_9 & -b_{10} & 0 & b_{11} \\ b_{10} & -1-b_9 & -b_{11} & 0 \end{pmatrix}$ $H_4 = \begin{pmatrix} 0 & b_{12} & b_{13} & -1-b_{14} \\ -b_{12} & 0 & b_{14} & b_{13} \\ -b_{13} & -b_{14} & 0 & b_{15} \\ 1+b_{14} & -b_{13} & -b_{15} & 0 \end{pmatrix}$
	$J = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$	$g = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$	$H_1 = \begin{pmatrix} 0 & c_1 & c_2 & c_3 \\ -c_1 & 0 & c_3 & c_4 \\ -c_2 & -c_3 & 0 & c_1 \\ -c_3 & -c_4 & -c_1 & 0 \end{pmatrix}$ $H_2 = \begin{pmatrix} 0 & c_5-1 & c_6 & c_7 \\ -c_5+1 & 0 & c_7 & c_8 \\ -c_6 & -c_7 & 0 & c_5 \\ -c_7 & -c_8 & -c_5 & 0 \end{pmatrix}$ $H_3 = \begin{pmatrix} 0 & c_9 & c_{10} & c_{11} \\ -c_9 & 0 & c_{11} & c_{12} \\ -c_{10} & -c_{11} & 0 & c_9 \\ -c_{11} & -c_{12} & -c_9 & 0 \end{pmatrix}$ $H_4 = \begin{pmatrix} 0 & c_{13} & c_{14} & c_{15} \\ -c_{13} & 0 & 1+c_{15} & c_{16} \\ -c_{14} & -1-c_{15} & 0 & c_{13} \\ -c_{15} & -c_{16} & -c_{13} & 0 \end{pmatrix}$

---

[6] Note that results of table 1 are solutions of the equations (9) and (12). But the results of table 2 are solutions of (9),(12) and (32),(33),(36),(39) so the results of table 2 can also be solutions of these equations which must be consistent with H and g .

TABLE 2 : biHermitian structures on four dimensional real Lie algebras

Lie Algebra	Complex Structures	g	antisymmetric tensor
$VIII \oplus R$	$J = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$	$g = \begin{pmatrix} -\alpha & 0 & 0 & 0 \\ 0 & -\alpha & 0 & 0 \\ 0 & 0 & \alpha & 0 \\ 0 & 0 & 0 & \alpha \end{pmatrix}$ $\alpha \in R - \{0\}$	$H_1 = \begin{pmatrix} 0 & d_1 & d_2 & -d_3+\alpha \\ -d_1 & 0 & d_3 & d_2 \\ -d_2 & -d_3 & 0 & d_4 \\ d_3-\alpha & -d_2 & -d_4 & 0 \end{pmatrix}$ $H_2 = \begin{pmatrix} 0 & d_5 & d_6 & -d_7 \\ -d_5 & 0 & d_7 & d_6+\alpha \\ -d_6 & -d_7 & 0 & d_8 \\ d_7 & -d_6-\alpha & -d_8 & 0 \end{pmatrix}$ $H_3 = \begin{pmatrix} 0 & d_9 & d_{10} & -d_{11} \\ -d_9 & 0 & d_{11} & d_{10} \\ -d_{10} & -d_{11} & 0 & d_{12} \\ d_{11} & -d_{10} & -d_{12} & 0 \end{pmatrix}$ $H_4 = \begin{pmatrix} 0 & d_{13} & d_{14} & -d_{15} \\ -d_{13} & 0 & d_{15} & d_{14} \\ -d_{14} & -d_{15} & 0 & d_{16} \\ d_{15} & -d_{14} & -d_{16} & 0 \end{pmatrix}$
$IX \oplus R$	$J = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$	$g = \begin{pmatrix} \beta & 0 & 0 & 0 \\ 0 & \beta & 0 & 0 \\ 0 & 0 & \beta & 0 \\ 0 & 0 & 0 & \beta \end{pmatrix}$ $\beta \in R - \{0\}$	$H_1 = \begin{pmatrix} 0 & 0 & f_1 & -f_2-\beta \\ 0 & 0 & f_2 & f_1 \\ -f_1 & -f_2 & 0 & f_3 \\ f_2+\beta & -f_1 & -f_3 & 0 \end{pmatrix}$ $H_2 = \begin{pmatrix} 0 & f_4 & f_5 & -f_6 \\ -f_4 & 0 & f_6 & f_5-\beta \\ -f_5 & -f_6 & 0 & f_7 \\ f_6 & -f_5+\beta & -f_7 & 0 \end{pmatrix}$ $H_3 = \begin{pmatrix} 0 & f_8 & f_9 & -f_{10} \\ -f_8 & 0 & f_{10} & f_9 \\ -f_9 & -f_{10} & 0 & f_{11} \\ f_{10} & -f_9 & -f_{11} & 0 \end{pmatrix}$ $H_4 = \begin{pmatrix} 0 & f_{12} & f_{13} & -f_{14} \\ -f_{12} & 0 & f_{14} & f_{13} \\ -f_{13} & -f_{14} & 0 & f_{15} \\ f_{14} & -f_{13} & -f_{15} & 0 \end{pmatrix}$

Note that b_i, c_i, d_i, f_i are all real parameters.

5 Conclusion

We offered a new method for calculation of complex and biHermitian structures on low dimensional Lie algebras. By this method, we obtain complex and biHermitian structures on real four dimensional Lie algebras. In this manner, one can obtain these structures on Lie groups using vierbeins . Some biHermitian structures on real four dimensional Lie algebras are equivalent to Manin triple structure obtained in [14]. One can use these methods for obtaining complex and biHermitian structures on real six dimensional Lie algebras [15]. We also apply this method for calculation of generalized complex structures on four dimensional real Lie algebras [16].

Acknowledgments

We would like to thank F. Darabi, Sh. Mogadassi and Z. Haghpanah for carefully reading the manuscript and useful comments.

Appendix

Real four dimensional Lie algebras and their automorphisms groups [9],[10]

TABLE A: Classifications of four dimensional real Lie algebras

Lie Algebra	Non Vanishing Structure Constants	Automorphisms group
$4A_1$		
$III \oplus R \cong (A_2 \oplus 2A_1)$	$f_{12}^2 = -1, f_{12}^3 = -1$ $, f_{31}^2 = 1, f_{31}^3 = 1$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_5 & a_4 & -a_6 \\ 0 & a_4 & a_5 & a_6 \\ 0 & a_7 & -a_7 & a_8 \end{pmatrix}$
$2A_2$	$f_{12}^2 = 1, f_{34}^4 = 1$	$\begin{pmatrix} 1 & a_1 & 0 & 0 \\ 0 & a_2 & 0 & 0 \\ 0 & 0 & 1 & a_3 \\ 0 & 0 & 0 & a_4 \end{pmatrix}$
$II \oplus R \cong (A_{3,1} \oplus A_1)$	$f_{23}^1 = 1$	$\begin{pmatrix} a_2 a_7 - a_3 a_6 & 0 & 0 & 0 \\ a_1 & a_2 & a_3 & a_4 \\ a_5 & a_6 & a_7 & a_8 \\ a_9 & 0 & 0 & a_{10} \end{pmatrix}$
$IV \oplus R \cong (A_{3,2} \oplus A_1)$	$f_{12}^2 = -1, f_{12}^3 = 1, f_{13}^3 = -1$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_4 & a_5 & 0 \\ 0 & 0 & a_4 & 0 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$
$V \oplus R \cong (A_{3,3} \oplus A_1)$	$f_{12}^2 = -1, f_{13}^3 = -1$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_4 & a_5 & 0 \\ 0 & a_6 & a_7 & 0 \\ 0 & 0 & 0 & a_8 \end{pmatrix}$
$VI_0 \oplus R \cong (A_{3,4} \oplus A_1)$	$f_{13}^2 = 1, f_{23}^1 = 1$	$\begin{pmatrix} a_2 & a_1 & 0 & 0 \\ a_1 & a_2 & 0 & 0 \\ a_3 & a_4 & 1 & a_5 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$
$VI_a \oplus R \cong \left(A_{3,5}^a \oplus A_1\right)$	$f_{12}^2 = -a, f_{12}^3 = -1$ $, f_{31}^2 = 1, f_{31}^3 = a$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_5 & a_4 & 0 \\ 0 & a_4 & a_5 & 0 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$
$VII_0 \oplus R \cong (A_{3,6} \oplus A_1)$	$f_{23}^1 = 1, f_{13}^2 = -1$	$\begin{pmatrix} a_2 & -a_1 & 0 & 0 \\ a_1 & a_2 & 0 & 0 \\ a_3 & a_4 & 1 & a_5 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$
$VII_a \oplus R \cong \left(A_{3,7}^a \oplus A_1\right)$	$f_{31}^2 = 1, f_{31}^3 = a$ $, f_{12}^2 = -a, f_{12}^3 = 1$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_5 & -a_4 & 0 \\ 0 & a_4 & a_5 & 0 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$

TABLE A: Classifications of four dimensional real Lie algebras

Lie Algebra	Non Vanishing Structure Constants	Automorphisms group
$VIII \oplus R \cong (A_{3,8} \oplus A_1)$	$f_{31}^2 = 1, f_{12}^3 = -1, f_{23}^1 = 1$	Λ_1
$IX \oplus R \cong (A_{3,9} \oplus A_1)$	$f_{31}^2 = 1, f_{12}^3 = 1, f_{23}^1 = 1$	Λ_2
$A_{4,1}$	$f_{24}^1 = 1, f_{34}^2 = 1$	$\begin{pmatrix} a_7^2 a_3 & 0 & 0 & 0 \\ a_2 a_7 & a_3 a_7 & 0 & 0 \\ a_1 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & a_7 \end{pmatrix}$
$A_{4,2}^a$	$f_{14}^1 = a, f_{24}^2 = 1$ $, f_{34}^2 = 1, f_{34}^3 = 1$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_3 & 0 & 0 \\ 0 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,2}^1$	$f_{14}^1 = 1, f_{24}^2 = 1$ $, f_{34}^2 = 1, f_{34}^3 = 1$	$\begin{pmatrix} a_1 & a_2 & 0 & 0 \\ 0 & a_5 & 0 & 0 \\ a_3 & a_4 & a_5 & 0 \\ a_6 & a_7 & a_8 & 1 \end{pmatrix}$
$A_{4,3}$	$f_{14}^1 = 1, f_{34}^2 = 1$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_2 & 0 & 0 \\ 0 & a_3 & a_2 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,4}$	$f_{14}^1 = 1, f_{24}^1 = 1, f_{24}^2 = 1$ $, f_{34}^2 = 1, f_{34}^3 = 1$	$\begin{pmatrix} a_3 & 0 & 0 & 0 \\ a_2 & a_3 & 0 & 0 \\ a_1 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,5}^{a,b}$	$f_{14}^1 = 1, f_{24}^2 = a$ $, f_{34}^3 = b$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_2 & 0 & 0 \\ 0 & 0 & a_3 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,5}^{a,a}$	$f_{14}^1 = 1, f_{24}^2 = a$ $, f_{34}^3 = a$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_2 & a_3 & 0 \\ 0 & a_4 & a_5 & 0 \\ a_6 & a_7 & a_8 & 1 \end{pmatrix}$
$A_{4,5}^{a,1}$	$f_{14}^1 = 1, f_{24}^2 = a$ $, f_{34}^3 = 1$	$\begin{pmatrix} a_1 & 0 & a_2 & 0 \\ 0 & a_3 & 0 & 0 \\ a_4 & 0 & a_5 & 0 \\ a_6 & a_7 & a_8 & 1 \end{pmatrix}$
$A_{4,5}^{1,1}$	$f_{14}^1 = 1, f_{24}^2 = 1$ $, f_{34}^3 = 1$	$\begin{pmatrix} a_1 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & 0 \\ a_7 & a_8 & a_9 & 0 \\ a_{10} & a_{11} & a_{12} & 1 \end{pmatrix}$

TABLE A: Classifications of four dimensional real Lie algebras

Lie Algebra	Non Vanishing Structure Constants	Automorphisms group
$A_{4,6}^{a,b}$	$f_{14}^1 = a, f_{24}^2 = b, f_{24}^3 = -1 , f_{34}^2 = 1, f_{34}^3 = b$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_3 & -a_2 & 0 \\ 0 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,7}$	$f_{14}^1 = 2, f_{24}^2 = 1, f_{34}^2 = 1 , f_{34}^3 = 1, f_{23}^1 = 1$	$\begin{pmatrix} a_2^2 & 0 & 0 & 0 \\ -a_2a_5 & a_2 & 0 & 0 \\ -a_2a_5 + a_2a_4 - a_1a_5 & a_1 & a_2 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,8}$	$f_{24}^2 = 1, f_{34}^3 = -1 , f_{23}^1 = 1$	$\begin{pmatrix} a_1a_2 & 0 & 0 & 0 \\ a_1a_5 & a_1 & 0 & 0 \\ a_2a_4 & 0 & a_2 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,9}^b$	$f_{14}^1 = 1 + b, f_{24}^2 = 1 , f_{34}^3 = b, f_{23}^1 = 1$	$\begin{pmatrix} a_1a_2 & 0 & 0 & 0 \\ -a_1a_5/b & a_1 & 0 & 0 \\ a_2a_4 & 0 & a_2 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,9}^1$	$f_{14}^1 = 2, f_{24}^2 = 1 , f_{34}^3 = 1, f_{23}^1 = 1$	$\begin{pmatrix} a_1a_4 - a_2a_3 & 0 & 0 & 0 \\ a_2a_6 - a_1a_7 & a_1 & a_2 & 0 \\ a_4a_6 - a_3a_7 & a_3 & a_4 & 0 \\ a_5 & a_6 & a_7 & 1 \end{pmatrix}$
$A_{4,9}^0$	$f_{14}^1 = 1, f_{24}^2 = 1 , f_{23}^1 = 1$	$\begin{pmatrix} a_2a_3 & 0 & 0 & 0 \\ a_1 & a_2 & 0 & 0 \\ a_3a_5 & 0 & a_3 & 0 \\ a_4 & a_5 & 0 & 1 \end{pmatrix}$
$A_{4,10}$	$f_{24}^3 = -1, f_{34}^2 = 1 , f_{23}^1 = 1$	$\begin{pmatrix} a_1^2 + a_2^2 & 0 & 0 & 0 \\ -a_1a_4 - a_2a_5 & a_1 & a_2 & 0 \\ a_2a_4 - a_1a_5 & -a_2 & a_1 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,11}^a$	$f_{14}^1 = 2a, f_{24}^2 = a, f_{24}^3 = -1 , f_{34}^2 = 1, f_{34}^3 = a, f_{23}^1 = 1$	$\begin{pmatrix} a_1^2 + a_2^2 & 0 & 0 & 0 \\ -\frac{a(a_1a_4)+a(a_2a_5)+a_2a_4-a_1a_5}{a^2+1} & a_2 & -a_1 & 0 \\ \frac{a(a_2a_4)-a(a_1a_5)-a_1a_4-a_2a_5}{a^2+1} & a_1 & a_2 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,12}$	$f_{14}^2 = -1, f_{13}^1 = 1 , f_{24}^1 = 1, f_{23}^2 = 1$	$\begin{pmatrix} a_2 & -a_1 & 0 & 0 \\ a_1 & a_2 & 0 & 0 \\ -a_4 & a_3 & 1 & 0 \\ a_3 & a_4 & 0 & 1 \end{pmatrix}$

$$\Lambda_1 = \text{Rotation}_{xy}\text{Boost}_{xz}\text{Boost}_{yz}C \tag{45}$$

where:

$$\text{Rotation}_{xy} = \begin{pmatrix} \cos(a_1) & \sin(a_1) & 0 & 0 \\ -\sin(a_1) & \cos(a_1) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{46}$$

$$\text{Boost}_{xz} = \begin{pmatrix} \cosh(a_2) & 0 & \sinh(a_2) & 0 \\ 0 & 1 & 0 & 0 \\ \sinh(a_2) & 0 & \cosh(a_2) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{47}$$

$$\text{Boost}_{yz} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cosh(a_3) & \sinh(a_3) & 0 \\ 0 & \sinh(a_3) & \cosh(a_3) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{48}$$

$$C = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & a_4 \end{pmatrix} \tag{49}$$

$$\tag{50}$$

$$\Lambda_2 = \text{Rotation}_{xy}\text{Rotation}_{xz}\text{Rotation}_{yz}C \tag{51}$$

where:

$$\text{Rotation}_{xy} = \begin{pmatrix} \cos(a_1) & \sin(a_1) & 0 & 0 \\ -\sin(a_1) & \cos(a_1) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{52}$$

$$\text{Rotation}_{xz} = \begin{pmatrix} \cos(a_2) & 0 & -\sin(a_2) & 0 \\ 0 & 1 & 0 & 0 \\ \sin(a_2) & 0 & \cos(a_2) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{53}$$

$$\text{Rotation}_{yz} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos(a_3) & \sin(a_3) & 0 \\ 0 & -\sin(a_3) & \cos(a_3) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{54}$$

$$C = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & a_4 \end{pmatrix} \tag{55}$$

References

[1] S.J.Gate, C.M.Hull and M.Roček,”*Twisted Multiplets and New Supersymmetric Nonlinear Sigma Model*”, Nucl.Phys. B **248**(1984) 157.

[2] S.Lyakhovich and M.Zabzine, ”*Poisson geometry of sigma models with extended supersymmetry* ”, Phys.Lett. B **548**(2002) 248, [arXiv:hep-th/0210043].

[3] S. Parkhomenko, ” *Extended superconformal current algebras and finite-dimensional Manin triples*”, Sov. Phys. JETP. 7 (1992),1-3.

[4] U.Lindström and M.Zabzine ,”*D-branes in N=2 WZW models*”, arXiv:hep-th/0212042 v2 5 Mar 2003.

[5] H.Samelson,” *A class of complex-analytic manifolds*”, Port.Math.(1953), 129-132.

[6] A.Morimoto,” *Structures complexes sur les groups de Lie semi-simples* ”, C.R.**242** (1956), 1101-1103.

[7] J.E.Snow ,”*Invariant Complex Structures on Four Dimensional Solvable Real Lie Groups*”, manuscripta math. **66**,(1990) 397-412 .

[8] G.Ovando ,”*Complex,Symplectic and Kähler structures on four dimensional Lie group*”,math.DG/0309146 v1,8 sep 2003,
”*Invariant complex structures on solvable real Lie groups*”, manuscripta math. **103**,(2000) 19-30 .

[9] J.Patera and P.Winternitz,” *Subalgebras of real three- and four dimensional Lie algebras*”, J.Math.Phys.**18**(7) (1977), 1449-1456.

[10] T.Christodoulakis, G.O.Papadopoulos and A.Dimakis ,”*Automorphisms of Real 4 Dimensional Lie Algebras and the Invariant Characterization of Homogeneous 4-Spaces*”, arXiv:gr-qc/0209042 v2 13 Nov 2002.

[11] A.Newlander and L.Nirenberg .Ann .Math **65** (1957) 391.

[12] A.Andrada, M.L.Barberis, I.G.Dotti, and G.P.Ovando ,”*Product structures on four dimensional solvable Lie algebras*”, math/0402234 v1 [math.RA],13 Feb 2004.

[13] S.E.Hjelmeland and U.Lindström,” *Comments on Dualities on Group Manifolds and Supersymmetry*”, arXiv:hep-th/0307137 v2 14 Aug 2003.

[14] L.Hlavaty and L.Snobl,” *Classification of Poisson-Lie T-dual models with two-dimensional targets*”, arXiv:hep-th/0110139 v2 20 Dec 2001.

[15] A. Rezaei-Aghdam and M. Sephid,” *Complex and biHermitian structures on six dimensional real Lie algebras*”. work in progress .

[16] S. Fallahpour, A. Rezaei-Aghdam , and S. Dadfar ” *Generelized complex structures on four dimensional real Lie algebras*”. work in progress .

arXiv:1002.4285v2 [math-ph] 13 Jul 2010

Complex and biHermitian structures on four dimensional real Lie algebras

A. Rezaei-Aghdam* and M. Sephid†
Department of Physics, Faculty of science, Azarbaijan University of Tarbiat Moallem , 53714-161, Tabriz, Iran

July 14, 2010

Abstract

We give a new method for calculation of complex and biHermitian structures on low dimensional real Lie algebras. In this method by use of non-coordinate basis, we first transform the Nijenhuis tensor field and biHermitian structure relations on Lie groups to the tensor relations on their Lie algebras. Then we use adjoint representation for writing these relations in the matrix form and by solving these matrix relations and use of automorphism groups of four dimensional real Lie algebras we obtain and classify all complex and biHermitian structures on four dimensional real Lie algebras.

---

*e-mail: rezaei-a@azaruniv.edu
†e-mail: s.sephid@azaruniv.edu

1 Introduction

Calculation of complex structures on homogeneous complex manifolds and especially on Lie groups are important from both mathematical and physical point of view. From mathematical discussion the classification of these manifolds are based on the determination of the possible complex structures. From the physical point of view these structures have important role in the N=(2,2) supersymmetric sigma models [1]. Its shown that the N=(2,2) extended supersymmetry in sigma model is equivalent to the existence of biHermitian structure on the target manifold such that the complex structures are covariantly constant with respect to torsionful affine connections (see for example [2] and references therein).Furthermore it is shown that the algebraic structures related to these bihermitian structure for N=(2,2) supersymmetric WZW models are the Manin triples [3] [4]. For these reasons the calculation of the complex and the biHermitian structures on manifolds especially on Lie groups are important. Samelson [5] shows that compact Lie groups always admit an invariant complex structures . In the non-compact case, Morimoto [6] showed that there always exist invariant complex structures on any even dimensional reductive Lie groups. In [7] and [8] complex structure on real four dimensional Lie algebra are classified. The method that used in those articles was special and seems not to be adequate for the calculation in higher dimensions . Here we give a new method for this purposes, which can be applied for low dimensional Lie groups . In this method, by use of non coordinate basis we first transform the Nijenhuis tensor on Lie groups to algebraic tensor relations on their Lie algebras. Then by use of adjoint representation we rewrite these relations in the matrix form. Finally, we solve these matrix relations by use of maple program. Here we perform this work for real four dimensional Lie algebras. Our consequences for some algebra are different and complete with respect to [8]. Furthermore, the calculation of biHermitian structure for four dimensional Lie algebras are new . The paper is organized as follows.

In section 2, by use of non coordinate basis we transform the Nijenhuis tensor relation to the algebraic tensor relation on Lie groups. Then by use of adjoint representation we write these relations in the matrix form. These relations can also be obtained from definition of the complex structures on Lie algebras as endomorphism of them. Then, in section 3 by use of maple program we solve these matrix relations to obtain the complex structures on real four dimensional Lie algebras. In this process, we apply automorphism groups of real four dimensional Lie algebras for obtaining of non equivalent complex structures (table 1). We then compare our results with [7] and [8]. Note that here we use Patera *etal* [9] classification of real four dimensional Lie algebras. The list of Lie algebras and their automorphism groups [10] are given in appendix. In section 4, we first transform the tensorial form of the biHermitian relations on Lie groups into the algebraic tensorial relations on their Lie algebras. In this respect, we define biHermitian structure on Lie algebra independently and give a proposition for obtaining non equivalent biHermitian structures. We then by use of adjoint representation rewrite these relations in the matrix forms and solve them by use of maple program. Our results for biHermitian structure on real four dimensional Lie algebra are new. Some discussions are given in the conclusion.

2 A brief review of complex structures on Lie groups

Definition 1: *Let M be a differentiable manifold, then the pair (M,J) is called almost complex manifold if there exists a tensor field J of type (1,1) such that at each point p of M , $J_p^2 = -1$. Tensor field J is also called the almost complex structure.*

Theorem : *An almost complex structure J on a manifold M is integrable if and only if*

$$N(X,Y) = 0 \qquad \forall X, Y \in \chi(M) \,, \tag{1}$$

where $\chi(M)$ is the set of vector fields on M and the Nijenhuis tensor N(X,Y) is given by

$$N(X,Y) = [X,Y] + J[JX,Y] + J[X,JY] - [JX,JY] \,. \tag{2}$$

In the coordinate basis i.e the basis $\{e_\mu = \frac{\partial}{\partial x^\mu}\}$ and $\{dx^\mu\}$ for vectors and dual vectors (forms) respectively on M the almost complex structures is expressed as $J = J_\mu{}^\nu e_\nu \otimes dx^\mu$ and the integrability condition can be rewritten as follows :

$$N(X,Y) = X^\kappa Y^\nu [J_\lambda{}^\mu(\partial_\kappa J_\nu{}^\lambda) + J_\nu{}^\lambda(\partial_\lambda J_\kappa{}^\mu) - J_\lambda{}^\mu(\partial_\nu J_\kappa{}^\lambda) - J_\kappa{}^\lambda(\partial_\lambda J_\nu{}^\mu)] e_\mu = 0. \tag{3}$$

Meanwhile the relation $J^2 = -1$ can be rewritten as

$$J_\lambda{}^\mu J_\mu{}^\nu = -\delta_\lambda{}^\nu. \tag{4}$$

Furthermore one can rewrite the above equations by use of the non-coordinate basis $\{\hat{e}_\alpha\}$ and $\{\hat{\theta}^\alpha\}$ for vectors and forms on M. For these basis we have

$$\hat{e}_\alpha = e_\alpha{}^\mu e_\mu \qquad , \qquad e_\alpha{}^\mu \in GL(m, R) \ , \tag{5}$$

where by requiring that $\{\hat{e}_\alpha\}$ be orthogonal we have the following relation for metric on M:

$$g_{\alpha\beta} = g(\hat{e}_\alpha, \hat{e}_\beta) = e_\alpha{}^\mu e_\beta{}^\nu g_{\mu\nu} \ , \tag{6}$$

or

$$g_{\mu\nu} = e^\alpha{}_\mu e^\beta{}_\nu g_{\alpha\beta} \ , \tag{7}$$

where $e^\alpha{}_\mu$ is the inverse of vielbein $e_\alpha{}^\mu$

$$e^\alpha{}_\mu e_\beta{}^\mu = \delta^\alpha{}_\beta \ \ , \ \ e^\alpha{}_\mu e_\alpha{}^\nu = \delta_\mu{}^\nu \ . \tag{8}$$

The dual basis $\{\hat{\theta^\alpha}\}$ are defined by $< \hat{\theta^\alpha}, \hat{e}_\beta >= \delta^\alpha{}_\beta$ and we have $\hat{\theta^\alpha} = e^\alpha{}_\mu dx^\mu$. Furthermore the vielbeins satisfy in the following relation:

$$f_{\alpha\beta}{}^\gamma = e^\gamma{}_\nu (e_\alpha{}^\mu \partial_\mu e_\beta{}^\nu - e_\beta{}^\mu \partial_\mu e_\alpha{}^\nu) \ , \tag{9}$$

where in the case M is a Lie group manifold G the $f_{\alpha\beta}{}^\gamma$,s are structure constants of Lie algebra **g** of G. Now by use of these basis the tensor $J = J_\mu{}^\nu e_\nu \otimes dx^\mu$ can be rewritten as

$$J_\mu{}^\nu = e^\alpha{}_\mu J_\alpha{}^\beta e_\beta{}^\nu \ , \tag{10}$$

where $J_\alpha{}^\beta$ is an endomorphism of **g** i.e. $J : \mathbf{g} \longrightarrow \mathbf{g}$. Now by applying this relation in (4) we have the following matrix relations for matrices $J_\alpha{}^\beta$:

$$J^2 = -I \ . \tag{11}$$

Furthermore by applying relations (9) and (10) in tensor equation (3) and use of (8) and assuming that $J_\alpha{}^\beta$ and $g_{\alpha\beta}$ are independent of coordinates of G, after some calculations we have the following algebraic relation for (3) :

$$f_{\beta\alpha}{}^\gamma + J_\sigma{}^\gamma \ J_\alpha{}^\delta \ f_{\beta\delta}{}^\sigma - J_\beta{}^\sigma \ J_\alpha{}^\delta \ f_{\sigma\delta}{}^\gamma + J_\beta{}^\sigma \ J_\delta{}^\gamma \ f_{\sigma\alpha}{}^\delta = 0. \tag{12}$$

Finally by using of adjoint representations

$$f_{\beta\alpha}{}^\gamma = -(\mathcal{Y}^\gamma)_{\beta\alpha} \quad , \quad f_{\beta\alpha}{}^\gamma = -(\chi_\beta)_\alpha{}^\gamma, \tag{13}$$

these relations have the following matrix forms:

$$\mathcal{Y}^\alpha + J \ \mathcal{Y}^\beta \ J_\beta{}^\alpha + J_\beta{}^\alpha \ \mathcal{Y}^\beta \ J^t - J \ \mathcal{Y}^\alpha \ J^t = 0 \ , \tag{14}$$

or

$$\chi_\alpha + J \ \chi_\alpha \ J + J_\alpha{}^\beta \ \chi_\beta \ J - J_\alpha{}^\beta \ J \ \chi_\beta = 0 \ . \tag{15}$$

Note that the above equation can also be obtained from the definition of complex structure on Lie algebra **g** as follows.

Definition 2: *An invariant complex structure on a real Lie algebra* **g** *is an endomorphism J of* **g** *such that*

a) $J^2 = -Id$, (16)

b) $[X, Y] + J[JX, Y] + J[X, JY] - [JX, JY] = 0 \ , \qquad \forall X, Y \in \mathbf{g}$. (17)

Now if we apply $\{X_\alpha\}$ as basis for Lie algebra **g** with the following structure constants:

$$[X_\alpha, X_\beta] = f_{\alpha\beta}{}^\gamma X_\gamma \ , \tag{18}$$

and use of the following relation for J:

$$J X_\alpha = J_\alpha{}^\beta X_\beta \ , \tag{19}$$

then relations (16) and (17) are rewritten as (11), (14) or (15). Now to obtain algebraic complex structures $J_\alpha{}^\beta$ it is enough to solve equations (11) and (14) or (15) simultaneously . We perform this work for real four dimensional Lie algebras in the next section.

3 Calculation of complex structures on four dimensional real Lie algebras

Here we use the Patera *etal* classification [9] of four dimensional real Lie algebras . The commutation relations and the automorphism groups of these Lie algebras are given in the appendix [1]. Now one can write the adjoint representation ($i.e\ \mathcal{Y}$) for these Lie algebras and then solve the matrix relations (11) and (14) for obtaining complex structures. We perform this work by use of maple program. Note that in this process one can obtain equivalent complex structures; to avoid these and obtain inequivalent complex structure we use the following proposition:

Proposition 1[8] *: Two complex structures* J_1 *and* J_2 *of Lie algebra* $\mathbf{g}$ *are equivalent if there exist an element* A *of automorphism group of Lie algebra* $\mathbf{g}$ *(*$Aut(\mathbf{g})$*) such that:*

$$J_2 = A\ J_1\ A^{-1}\ . \tag{20}$$

In this way, we perform this work and obtain all nonequivalent complex structures on four dimensional real Lie algebras. The results are classified in the table 1 [2]. We see that 21 out of 30 real four dimensional Lie algebras have complex structures. To comparison of these results with the results of [8] we must firstly obtain the isomorphism relations between the four dimensional real Lie algebras presented in [9] and those presented in [8]. According to the calculations [8] and [11] we have isomorphism relations as the following table:

$4A_1$	$A_{4,1}$	$A^a_{4,2}$	$A_{4,3}$	$A_{4,4}$	$A^{a,b}_{4,5}$	$A^{a,b}_{4,6}$	$A_{4,7}$	$A_{4,8}$	$A^b_{4,9}$	$A_{4,10}$	$A^a_{4,11}$	$A_{4,12}$
$\mathfrak{a}_4$	$\mathfrak{n}_4$	$\mathfrak{r}_{4,a}$	$\mathfrak{r}_{4,0}$	$\mathfrak{r}_4$	$\mathfrak{r}_{4,a,b}$	$\mathfrak{r}'_{4,a,b}$	$\mathfrak{h}_4$	$\mathfrak{d}_4$	$\mathfrak{d}_{4,1/1+b}$	$\mathfrak{d}'_{4,0}$	$\mathfrak{d}'_{4,a}$	$\mathfrak{aff}(\mathbb{C})$

$A_2 \oplus A_2$	$II \oplus R$	$III \oplus R$	$IV \oplus R$	$V \oplus R$	$VI_0 \oplus R$	$VI_a \oplus R$ $(a \neq 0,1)$	$VII_0 \oplus R$	$VII_a \oplus R$ $(a \neq 0)$
$\mathfrak{r}_2\mathfrak{r}_2$	$\mathfrak{rh}_3$	$\mathfrak{rr}_{3,0}$	$\mathfrak{rr}_3$	$\mathfrak{rr}_{3,1}$	$\mathfrak{rr}_{3,-1}$	$\mathfrak{rr}_{3,a}$	$\mathfrak{rr}'_{3,0}$	$\mathfrak{rr}'_{3,a}$

In this respect one can see that in [8] for Lie algebra $VII_0 + R$ one complex structure is obtained but according to our calculation this Lie algebra has two non-equivalent complex structures.

For non solvable Lie algebras $VIII + R$ and $IX + R$ we obtain complex structures.

In [8], for Lie algebra $A_{4,8}$, two complex structures are obtained but here we obtain four complex structures to this Lie algebra.

For Lie algebra $A_{4,10}$ two complex structures are obtained in [8] but according to table 1,we obtain four complex structures for this Lie algebras.

In [8], for Lie algebra $A_{4,12}$, two complex structures are obtained but here we obtain three complex structures for this Lie algebra.

The results for other Lie algebras are the same as [8].

---

[1] Note that for decomposable Lie algebra ($L_3 \oplus R$)we use the Bianchi classification for real three dimensional Lie algebras L_3.

[2] Note that in this table the basis are showed with $\{e_\alpha\}$ instead $\{X_\alpha\}$

TABLE 1 : Complex structures on four dimensional real Lie algebras

Lie Algebra	Complex Structures
$III \oplus R$	$Je_1 = -e_2 \ , \ Je_3 = e_1 - e_4$
$A_2 \oplus A_2$	$Je_1 = -e_2 \ , \ Je_3 = -e_4$
$II \oplus R$	$Je_1 = -e_4 \ , \ Je_2 = e_3$
$V \oplus R$	$Je_1 = -e_4 \ , \ Je_2 = e_3$
$VII_0 \oplus R$	$Je_1 = -e_2 \ , \ Je_3 = -e_4$
	$Je_1 = e_2 \ , \ Je_3 = -e_4$
$VII_a \oplus R$ $(a \neq 0)$	$Je_1 = -e_4 \ , \ Je_2 = e_3$
	$Je_1 = -e_4 \ , \ Je_2 = -e_3$
$VIII \oplus R$	$Je_1 = -e_2 \ , \ Je_3 = pe_3 - (1 + p^2)e_4 \qquad (p \in \mathbb{R})$
$IX \oplus R$	$Je_1 = -e_2 \ , \ Je_3 = pe_3 - (1 + p^2)e_4 \qquad (p \in \mathbb{R})$
$A^1_{4,2}$	$Je_1 = e_2 \ , \ Je_3 = e_4$
$A^{a,a}_{4,5}$ $(-1 \leq a < 1, a \neq 0)$	$Je_1 = -e_4 \ , \ Je_2 = e_3$
$A^{a,1}_{4,5}$ $(-1 \leq a < 1, a \neq 0)$	$Je_1 = e_3 \ , \ Je_2 = e_4$
$A^{1,1}_{4,5}$	$Je_1 = -e_3 \ , \ Je_2 = -e_4$
$A^{a,b}_{4,6}$ $(a \neq 0, b \geq 0)$	$Je_1 = e_4 \ , \ Je_2 = e_3$
	$Je_1 = e_4 \ , \ Je_2 = -e_3$
$A_{4,7}$	$Je_1 = e_2 \ , \ Je_3 = -e_4$
$A_{4,8}$	$Je_1 = e_2 \ , \ Je_3 = e_4$
	$Je_1 = -e_3 \ , \ Je_2 = -e_4$
	$Je_1 = e_2 \ , \ Je_3 = e_2 + e_4$
	$Je_1 = -(e_1 + 2e_3) \ , \ Je_2 = -(e_1 + e_2) - 2(e_3 + e_4)$
$A^b_{4,9}$ $(0 <\mid b \mid< 1)$	$Je_1 = -be_2 \ , \ Je_3 = e_4$
	$Je_1 = -e_3 \ , \ Je_2 = -e_4$
$A^1_{4,9}$	$Je_1 = -e_3 \ , \ Je_2 = -e_4$
	$Je_1 = e_4 \ , \ Je_2 = -e_3$
	$Je_1 = -e_4 \ , \ Je_2 = -e_3$
$A^0_{4,9}$	$Je_1 = -e_3 \ , \ Je_2 = -e_4$
$A_{4,10}$	$Je_2 = -e_1 + e_3 \ , \ Je_3 = -e_1 - e_2 + e_3 + e_4$
	$Je_2 = e_1 + e_3 \ , \ Je_3 = -e_1 - e_2 - e_3 + e_4$
	$Je_2 = -e_1 - e_3 \ , \ Je_3 = e_1 + e_2 + e_3 - e_4$
	$Je_2 = -e_1 - e_3 \ , \ Je_3 = -e_1 + e_2 - e_3 + e_4$
$A^a_{4,11}$ $(a > 0)$	$Je_1 = e_4 \ , \ Je_2 = -e_3$
	$Je_1 = e_4 \ , \ Je_2 = e_3$
	$Je_1 = -e_4 \ , \ Je_2 = -e_3$
	$Je_1 = -e_4 \ , \ Je_2 = e_3$
$A_{4,12}$	$Je_1 = e_2 \ , \ Je_3 = e_4$
	$Je_1 = -e_2 \ , \ Je_3 = e_4$
	$Je_1 = e_4 \ , \ Je_2 = e_3$

4 BiHermitian structures on four dimensional real Lie algebras

Definition 3 [1] [2] :*If the manifold M has two complex structures $J_{\pm}$ such that these are biHermitian ,i.e Hermitian with respect to both complex structures*

$$J_{\pm}^{2} = -1 \ , \tag{21}$$

$$N_{\mu\nu}{}^{\kappa}(J_{\pm}) = 0 \ , \tag{22}$$

$$J_{\pm\mu}{}^{\lambda}\ g_{\lambda\eta}\ J_{\pm\nu}{}^{\eta} = g_{\mu\nu}, \tag{23}$$

where $g_{\mu\nu}$ is the metric on M , and furthermore these complex structures be covariantly constant with respect to certain connections $\Gamma^{\pm}$, respectively

$$\bigtriangledown_{\mu}^{\pm} \quad J_{\pm\nu}{}^{\lambda} \equiv J_{\pm\nu}{}^{\lambda}{}_{,\mu} + \Gamma^{\pm\ \lambda}_{\mu\rho} J_{\pm\nu}{}^{\rho} - \Gamma^{\pm\ \rho}_{\mu\nu} J_{\pm\rho}{}^{\lambda} = 0, \tag{24}$$

such that

$$\Gamma^{\pm\ \lambda}_{\mu\nu} = \Gamma_{\mu\nu}{}^{\lambda} \pm T_{\mu\nu}{}^{\lambda}, \tag{25}$$

where $\Gamma_{\mu\nu}{}^{\lambda}$ is the Christofel connection for the metric g and the torsion is given by

$$T_{\mu\nu}{}^{\lambda} = H_{\mu\nu\eta} g^{\eta\lambda}, \tag{26}$$

so that $H_{\mu\nu\eta}$ is antisymmetric tensor ; then it is said that M has biHermitian structure and is shown by $(M, g, J_{\pm})$.

Using (24) the integrebility condition (22) may be rewritten in the following form [2]:

$$H_{\delta\nu\lambda} = J_{\pm\delta}{}^{\sigma} J_{\pm\nu}{}^{\rho} H_{\sigma\rho\lambda} + J_{\pm\delta}{}^{\rho} J_{\pm\lambda}{}^{\sigma} H_{\sigma\rho\nu} + J_{\pm\upsilon}{}^{\sigma} J_{\pm\lambda}{}^{\rho} H_{\sigma\rho\delta}\ . \tag{27}$$

Furthermore by introducing the Kähler forms

$$\omega_{\pm\mu\nu} \equiv\ g_{\mu\nu} J_{\pm\nu}{}^{\lambda}, \tag{28}$$

and by use of (24) one can find

$$(d\omega_{\pm})_{\rho\mu\nu} = \pm(H_{\sigma\rho\mu} J^{\sigma}_{\pm\nu} + H_{\sigma\mu\nu} J^{\sigma}_{\pm\rho} + H_{\sigma\nu\rho} J^{\sigma}_{\pm\mu}), \tag{29}$$

where

$$(d\omega_{\pm})_{\lambda\sigma\gamma} = \frac{1}{2}(\partial_{\lambda}\omega_{\pm\sigma\gamma} + \partial_{\sigma}\omega_{\pm\gamma\lambda} + \partial_{\gamma}\omega_{\pm\lambda\sigma}). \tag{30}$$

Finally by use of (28) and (29) one can find

$$H_{\mu\nu\rho} = -J^{\lambda}_{+\mu} J^{\sigma}_{+\nu} J^{\gamma}_{+\rho} (d\omega_{+})_{\lambda\sigma\gamma} = -J^{\lambda}_{-\mu} J^{\sigma}_{-\nu} J^{\gamma}_{-\rho} (d\omega_{-})_{\lambda\sigma\gamma}. \tag{31}$$

In this respect, the target manifold $(M, g, J_{\pm})$ is said to have biHermitian structure if two Hermitian complex structures $J_{\pm}$ satisfy the relation (31) (i.e relation between (J_{+}, ω_{+}) and (J_{-}, ω_{-})) defining the torsion H. Now for the case where M is a Lie group G in the similar way of section 2 one can transform relations (21) – (24) and (27) to the algebric relation by using the relations (7) – (9) and the following relations:

$$H_{\mu\nu\rho} = \frac{1}{2} L^{\alpha}{}_{\mu} L^{\beta}{}_{\nu} L^{\gamma}{}_{\rho} f_{\alpha\beta\gamma} = \frac{1}{2} R^{\alpha}{}_{\mu} R^{\beta}{}_{\nu} R^{\gamma}{}_{\rho} f_{\alpha\beta\gamma}, \tag{32}$$

$$J_{+\mu}{}^{\nu} = R^{\alpha}{}_{\mu} J_{\alpha}{}^{\beta} R_{\beta}{}^{\nu}\ ,\ J_{-\mu}{}^{\nu} = L^{\alpha}{}_{\mu} J_{\alpha}{}^{\beta} L_{\beta}{}^{\nu}, \tag{33}$$

where $L^{\alpha}{}_{\mu}(R^{\alpha}{}_{\mu})$ and $L_{\beta}{}^{\nu}(R_{\beta}{}^{\nu})$ are left(right) invariant vielbeins and their inverses respectively. Now by using these relations , (23) and (27) transform to the following matrix relations:

$$J\ g\ J^{t} = g, \tag{34}$$

$$H_{\alpha} = J(H_{\beta} J_{\alpha}{}^{\beta}) + J H_{\alpha} J^{t} + (H_{\beta} J_{\alpha}{}^{\beta}) J^{t}, \tag{35}$$

where $(H_\alpha)_{\beta\gamma} = H_{\alpha\beta\gamma}$. Furthermore by using the following relations: [3]

$$\bigtriangledown^\rho L_\alpha{}^\mu = -\frac{1}{2}(f_\alpha{}^{(\rho\mu)} + f^{\rho\mu}{}_\alpha + T_\alpha{}^{(\rho\mu)} + T^{\rho\mu}{}_\alpha + L_\beta{}^\rho L_\gamma{}^\mu \bigtriangledown_\alpha g^{\beta\gamma} + L^{\beta\rho} \bigtriangledown^\mu g_{\alpha\beta} - L^{\beta\mu} \bigtriangledown^\rho g_{\alpha\beta}), \qquad (36)$$

$$\bigtriangledown^\rho R_\alpha{}^\mu = -\frac{1}{2}(-f_\alpha{}^{(\rho\mu)} - f^{\rho\mu}{}_\alpha + T_\alpha{}^{(\rho\mu)} + T^{\rho\mu}{}_\alpha + R_\beta{}^\rho R_\gamma{}^\mu \bigtriangledown_\alpha g^{\beta\gamma} + R^{\beta\rho} \bigtriangledown^\mu g_{\alpha\beta} - R^{\beta\mu} \bigtriangledown^\rho g_{\alpha\beta}), \qquad (37)$$

and by assuming that $g_{\alpha\beta}$ are coordinate independent; the relation (24) transforms to the following algebraic relation[4]:

$$J(H_\alpha - \chi_\alpha g) = (J(H_\alpha - \chi_\alpha g))^t. \qquad (38)$$

Note that the metric $g_{\alpha\beta}$ is the ad invariant metric on Lie algebras **g** .i.e. we have

$$\langle X_\alpha, X_\beta \rangle = g_{\alpha\beta}, \qquad (39)$$

$$\langle X_\alpha, [X_\beta, X_\gamma] \rangle = \langle [X_\alpha, X_\beta], X_\gamma \rangle, \qquad (40)$$

or in matrix notation we have

$$\chi_\alpha g = -(\chi_\alpha g)^t. \qquad (41)$$

Now, one can obtain biHermitian structures on Lie algebras by solving relations (11), (14), (34), (35), (38) and (41) simultaneously. These relations can be applied as a definition of algebraic biHermitian structure on the Lie algebra **g** ;

Definition 4: *If there exist endomorphism $J : \mathbf{g} \rightarrow \mathbf{g}$ of Lie algebra with ad invariant metric g and anti-symmetric bilinear map $H : \mathbf{g} \otimes \mathbf{g} \rightarrow \mathbf{g}$ such that the relations* (11), (14), (34), (35), (38) *and* (41) *are satisfied, then we have biHermitian structure (J, g, H) on* **g**.

Note that relation (35) is equivalent to matrix relation of integrability condition i.e relation (14) . For this reason, first it is better to obtain algebraic complex structures J, then solve relations (34), (38) and (41) and finally check them in (35). We perform this work for real four dimensional Lie algebras by use of maple program. Note that as in the case of complex structures to obtain non equivalent biHermitian structures we propose the following proposition:

Proposition 2 *: Two biHermitian structures (J, g, H) and $(J^{'}, g^{'}, H^{'})$ of Lie algebras* **g** *are equivalent if there exist an element A of automorphism group of Lie algebra* **g***(Auto* **g***) such that:*

$$J^{'} = AJA^{-1}, \qquad (42)$$

$$g^{'} = AgA^t, \qquad (43)$$

$$H^{'}_\alpha = A(H_\beta A_\alpha{}^\beta)A^t. \qquad (44)$$

Proof: *By use of* (14), (34) *and* (35) *one can see that if (J, g, H) is a biHermitian structure then $(J^{'}, g^{'}, H^{'})$ is also biHermitian structure and satisfy in those relations.*

Note that in the case that $f_{\beta\gamma}{}^\alpha = H_{\delta\beta\gamma} g^{\delta\alpha}$ or H is isomorphic with f i.e if there exist isomorphism matrix C so that

$$CY^\alpha C^t = \tilde{Y}^\beta C_\beta{}^\alpha, \qquad (45)$$

where $(Y^\alpha)_{\beta\gamma} = -f_{\beta\gamma}{}^\alpha$ and $(\tilde{Y}^\alpha)_{\beta\gamma} = -H_{\delta\beta\gamma} g^{\delta\alpha}$; then (J, g, H) shows the Manin triple structure on **g**[4]. In this way biHermitian structures on real four dimensional Lie algebras are classified in the following table . Note that according to the table 2 for Lie algebra $A_{4,8}$ we have two nonequivalent biHermitian structure (J, g, H)

---

[3] Note that in these relations all algebraic (target) indices are lowered and raised by $g_{\alpha\beta}(g_{\mu\nu})$; furthermore these indices transform into each other by $L_\alpha{}^\mu(R_\alpha{}^\mu)$ or $L^\alpha{}_\mu(R^\alpha{}_\mu)$. The symmetrization notation have the following form: $f_\alpha{}^{(\rho\mu)} = f_\alpha{}^{\rho\mu} + f_\alpha{}^{\mu\rho}$.

[4] This relation can also be obtained from algebraic form of (31).

where the second biHermitian structure shows the Manin triple structure of $A_{4,8}$ [4] (i.e $A_{4,8}$ is a Manin triple of two dimensional Lie bialgebras (type B and semiabelian)[12]) for the following values of parameters:

$$c_1 = c_2 = c_3 = c_4 = c_5 = c_6 = c_7 = c_9 = c_{10} = c_{11} = c_{13} = c_{14} = 0\ ,\ c_{12} = c_{15} = -1.$$

For Lie algebras $VIII \oplus R$ there is one biHermitian structure where this structure for the values

$$d_1 = d_2 = d_4 = d_5 = d_7 = d_8 = d_{10} = d_{11} = d_{12} = d_{14} = d_{15} = d_{16} = 0\ ,\ d_9 \neq 0\ ,\ \ d_3 = -d_6 = \alpha,$$

is isomorphic with two dimensional Lie bialgebra type A [12]. There exist one biHermotian structure for Lie algebra $IX \oplus R$. The results are given in the table 2. Note that the isomorphism relation (45)(i.e. the biHermitian structures which shows Manin triple) are independent of the choice of special biHermitian structures from equivalent class of biHermitian structures. In this way if relation (45) holds ; then by use of $\tilde{Y}'^{\alpha} = -H'_{\delta} g'^{\delta\alpha}$ and relations (44) and (45) one can show that

$$(AC)Y^{\gamma}(AC)^t = \tilde{Y}'^{\alpha}(AC)_{\alpha}{}^{\gamma}. \qquad (46)$$

TABLE 2 : biHermitian structures on four dimensional real Lie algebras

Lie Algebra	Complex Structures	g	antisymmetric tensor
$A_{4,8}$	$J = \begin{pmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{pmatrix}$	$g = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$	$H_1 = \begin{pmatrix} 0 & b_1 & 1 & -b_2 \\ -b_1 & 0 & b_2 & 1 \\ -1 & -b_2 & 0 & b_3 \\ b_2 & -1 & -b_3 & 0 \end{pmatrix}$ $H_2 = \begin{pmatrix} 0 & b_4 & b_5 & -b_6 \\ -b_4 & 0 & b_6 & b_5 \\ -b_5 & -b_6 & 0 & b_7 \\ b_6 & -b_5 & -b_7 & 0 \end{pmatrix}$ $H_3 = \begin{pmatrix} 0 & b_8 & b_9 & -b_{10} \\ -b_8 & 0 & b_{10} & 1+b_9 \\ -b_9 & -b_{10} & 0 & b_{11} \\ b_{10} & -1-b_9 & -b_{11} & 0 \end{pmatrix}$ $H_4 = \begin{pmatrix} 0 & b_{12} & b_{13} & -1-b_{14} \\ -b_{12} & 0 & b_{14} & b_{13} \\ -b_{13} & -b_{14} & 0 & b_{15} \\ 1+b_{14} & -b_{13} & -b_{15} & 0 \end{pmatrix}$
	$J = \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}$	$g = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$	$H_1 = \begin{pmatrix} 0 & c_1 & c_2 & c_3 \\ -c_1 & 0 & c_3 & c_4 \\ -c_2 & -c_3 & 0 & c_1 \\ -c_3 & -c_4 & -c_1 & 0 \end{pmatrix}$ $H_2 = \begin{pmatrix} 0 & c_5-1 & c_6 & c_7 \\ -c_5+1 & 0 & c_7 & c_8 \\ -c_6 & -c_7 & 0 & c_5 \\ -c_7 & -c_8 & -c_5 & 0 \end{pmatrix}$ $H_3 = \begin{pmatrix} 0 & c_9 & c_{10} & c_{11} \\ -c_9 & 0 & c_{11} & c_{12} \\ -c_{10} & -c_{11} & 0 & c_9 \\ -c_{11} & -c_{12} & -c_9 & 0 \end{pmatrix}$ $H_4 = \begin{pmatrix} 0 & c_{13} & c_{14} & c_{15} \\ -c_{13} & 0 & 1+c_{15} & c_{16} \\ -c_{14} & -1-c_{15} & 0 & c_{13} \\ -c_{15} & -c_{16} & -c_{13} & 0 \end{pmatrix}$

TABLE 2 : biHermitian structures on four dimensional real Lie algebras

Lie Algebra	Complex Structures	g	antisymmetric tensor
$VIII \oplus R$	$J = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$	$g = \begin{pmatrix} -\alpha & 0 & 0 & 0 \\ 0 & -\alpha & 0 & 0 \\ 0 & 0 & \alpha & 0 \\ 0 & 0 & 0 & \alpha \end{pmatrix}$ $\alpha \in R - \{0\}$	$H_1 = \begin{pmatrix} 0 & d_1 & d_2 & -d_3+\alpha \\ -d_1 & 0 & d_3 & d_2 \\ -d_2 & -d_3 & 0 & d_4 \\ d_3-\alpha & -d_2 & -d_4 & 0 \end{pmatrix}$ $H_2 = \begin{pmatrix} 0 & d_5 & d_6 & -d_7 \\ -d_5 & 0 & d_7 & d_6+\alpha \\ -d_6 & -d_7 & 0 & d_8 \\ d_7 & -d_6-\alpha & -d_8 & 0 \end{pmatrix}$ $H_3 = \begin{pmatrix} 0 & d_9 & d_{10} & -d_{11} \\ -d_9 & 0 & d_{11} & d_{10} \\ -d_{10} & -d_{11} & 0 & d_{12} \\ d_{11} & -d_{10} & -d_{12} & 0 \end{pmatrix}$ $H_4 = \begin{pmatrix} 0 & d_{13} & d_{14} & -d_{15} \\ -d_{13} & 0 & d_{15} & d_{14} \\ -d_{14} & -d_{15} & 0 & d_{16} \\ d_{15} & -d_{14} & -d_{16} & 0 \end{pmatrix}$
$IX \oplus R$	$J = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$	$g = \begin{pmatrix} \beta & 0 & 0 & 0 \\ 0 & \beta & 0 & 0 \\ 0 & 0 & \beta & 0 \\ 0 & 0 & 0 & \beta \end{pmatrix}$ $\beta \in R - \{0\}$	$H_1 = \begin{pmatrix} 0 & 0 & f_1 & -f_2-\beta \\ 0 & 0 & f_2 & f_1 \\ -f_1 & -f_2 & 0 & f_3 \\ f_2+\beta & -f_1 & -f_3 & 0 \end{pmatrix}$ $H_2 = \begin{pmatrix} 0 & f_4 & f_5 & -f_6 \\ -f_4 & 0 & f_6 & f_5-\beta \\ -f_5 & -f_6 & 0 & f_7 \\ f_6 & -f_5+\beta & -f_7 & 0 \end{pmatrix}$ $H_3 = \begin{pmatrix} 0 & f_8 & f_9 & -f_{10} \\ -f_8 & 0 & f_{10} & f_9 \\ -f_9 & -f_{10} & 0 & f_{11} \\ f_{10} & -f_9 & -f_{11} & 0 \end{pmatrix}$ $H_4 = \begin{pmatrix} 0 & f_{12} & f_{13} & -f_{14} \\ -f_{12} & 0 & f_{14} & f_{13} \\ -f_{13} & -f_{14} & 0 & f_{15} \\ f_{14} & -f_{13} & -f_{15} & 0 \end{pmatrix}$

5 Conclusion

We give new method for calculation of complex and biHermitian structures on low dimensional Lie algebras. By use of this method we obtain complex and biHermitian structures on real four dimensional Lie algebras. In this manner one can obtain these structures on Lie groups by use of veilbins . Some biHermitian structure on real four dimensional Lie algebras are equivalent to Manin triple structure that have been obtained previously [12]. One can apply these methods for obtaining of complex and biHermitian structures on real six dimensional Lie algebras [13].

Acknowledgments

We would like to thank F. Darabi and Z. Haghpanah for carefully reading the manuscript and useful comments.

Appendix A

Real four dimensional Lie algebras and their automorphisms groups [9],[10]

TABLE A: Classifications of four dimensional real Lie algebras

Lie Algebra	Non Vanishing Structure Constants	Automorphisms group
$4A_1$		
$III \oplus R \cong (A_2 \oplus 2A_1)$	$f_{12}^2 = -1, f_{12}^3 = -1$ $, f_{31}^2 = 1, f_{31}^3 = 1$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_5 & a_4 & -a_6 \\ 0 & a_4 & a_5 & a_6 \\ 0 & a_7 & -a_7 & a_8 \end{pmatrix}$
$2A_2$	$f_{12}^2 = 1, f_{34}^4 = 1$	$\begin{pmatrix} 1 & a_1 & 0 & 0 \\ 0 & a_2 & 0 & 0 \\ 0 & 0 & 1 & a_3 \\ 0 & 0 & 0 & a_4 \end{pmatrix}$
$II \oplus R \cong (A_{3,1} \oplus A_1)$	$f_{23}^1 = 1$	$\begin{pmatrix} a_2a_7 - a_3a_6 & 0 & 0 & 0 \\ a_1 & a_2 & a_3 & a_4 \\ a_5 & a_6 & a_7 & a_8 \\ a_9 & 0 & 0 & a_{10} \end{pmatrix}$
$IV \oplus R \cong (A_{3,2} \oplus A_1)$	$f_{12}^2 = -1, f_{12}^3 = 1, f_{13}^3 = -1$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_4 & a_5 & 0 \\ 0 & 0 & a_4 & 0 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$
$V \oplus R \cong (A_{3,3} \oplus A_1)$	$f_{12}^2 = -1, f_{13}^3 = -1$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_4 & a_5 & 0 \\ 0 & a_6 & a_7 & 0 \\ 0 & 0 & 0 & a_8 \end{pmatrix}$
$VI_0 \oplus R \cong (A_{3,4} \oplus A_1)$	$f_{13}^2 = 1, f_{23}^1 = 1$	$\begin{pmatrix} a_2 & a_1 & 0 & 0 \\ a_1 & a_2 & 0 & 0 \\ a_3 & a_4 & 1 & a_5 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$
$VI_a \oplus R \cong \left(A_{3,5}^a \oplus A_1\right)$	$f_{12}^2 = -a, f_{12}^3 = -1$ $, f_{31}^2 = 1, f_{31}^3 = a$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_5 & a_4 & 0 \\ 0 & a_4 & a_5 & 0 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$
$VII_0 \oplus R \cong (A_{3,6} \oplus A_1)$	$f_{23}^1 = 1, f_{13}^2 = -1$	$\begin{pmatrix} a_2 & -a_1 & 0 & 0 \\ a_1 & a_2 & 0 & 0 \\ a_3 & a_4 & 1 & a_5 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$
$VII_a \oplus R \cong \left(A_{3,7}^a \oplus A_1\right)$	$f_{31}^2 = 1, f_{31}^3 = a$ $, f_{12}^2 = -a, f_{12}^3 = 1$	$\begin{pmatrix} 1 & a_1 & a_2 & a_3 \\ 0 & a_5 & -a_4 & 0 \\ 0 & a_4 & a_5 & 0 \\ 0 & 0 & 0 & a_6 \end{pmatrix}$

TABLE A: Classifications of four dimensional real Lie algebras

Lie Algebra	Non Vanishing Structure Constants	Automorphisms group
$VIII \oplus R \cong (A_{3,8} \oplus A_1)$	$f_{31}^2 = 1, f_{12}^3 = -1, f_{23}^1 = 1$	Λ_1
$IX \oplus R \cong (A_{3,9} \oplus A_1)$	$f_{31}^2 = 1, f_{12}^3 = 1, f_{23}^1 = 1$	Λ_2
$A_{4,1}$	$f_{24}^1 = 1, f_{34}^2 = 1$	$\begin{pmatrix} a_7^2 a_3 & 0 & 0 & 0 \\ a_2 a_7 & a_3 a_7 & 0 & 0 \\ a_1 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & a_7 \end{pmatrix}$
$A_{4,2}^a$	$f_{14}^1 = a, f_{24}^2 = 1$ $, f_{34}^2 = 1, f_{34}^3 = 1$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_3 & 0 & 0 \\ 0 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,2}^1$	$f_{14}^1 = 1, f_{24}^2 = 1$ $, f_{34}^2 = 1, f_{34}^3 = 1$	$\begin{pmatrix} a_1 & a_2 & 0 & 0 \\ 0 & a_5 & 0 & 0 \\ a_3 & a_4 & a_5 & 0 \\ a_6 & a_7 & a_8 & 1 \end{pmatrix}$
$A_{4,3}$	$f_{14}^1 = 1, f_{34}^2 = 1$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_2 & 0 & 0 \\ 0 & a_3 & a_2 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,4}$	$f_{14}^1 = 1, f_{24}^1 = 1, f_{24}^2 = 1$ $, f_{34}^2 = 1, f_{34}^3 = 1$	$\begin{pmatrix} a_3 & 0 & 0 & 0 \\ a_2 & a_3 & 0 & 0 \\ a_1 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,5}^{a,b}$	$f_{14}^1 = 1, f_{24}^2 = a$ $, f_{34}^3 = b$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_2 & 0 & 0 \\ 0 & 0 & a_3 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,5}^{a,a}$	$f_{14}^1 = 1, f_{24}^2 = a$ $, f_{34}^3 = a$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_2 & a_3 & 0 \\ 0 & a_4 & a_5 & 0 \\ a_6 & a_7 & a_8 & 1 \end{pmatrix}$
$A_{4,5}^{a,1}$	$f_{14}^1 = 1, f_{24}^2 = a$ $, f_{34}^3 = 1$	$\begin{pmatrix} a_1 & 0 & a_2 & 0 \\ 0 & a_3 & 0 & 0 \\ a_4 & 0 & a_5 & 0 \\ a_6 & a_7 & a_8 & 1 \end{pmatrix}$
$A_{4,5}^{1,1}$	$f_{14}^1 = 1, f_{24}^2 = 1$ $, f_{34}^3 = 1$	$\begin{pmatrix} a_1 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & 0 \\ a_7 & a_8 & a_9 & 0 \\ a_{10} & a_{11} & a_{12} & 1 \end{pmatrix}$

TABLE A: Classifications of four dimensional real Lie algebras

Lie Algebra	Non Vanishing Structure Constants	Automorphisms group
$A_{4,6}^{a,b}$	$f_{14}^1 = a, f_{24}^2 = b, f_{24}^3 = -1$ $, f_{34}^2 = 1, f_{34}^3 = b$	$\begin{pmatrix} a_1 & 0 & 0 & 0 \\ 0 & a_3 & -a_2 & 0 \\ 0 & a_2 & a_3 & 0 \\ a_4 & a_5 & a_6 & 1 \end{pmatrix}$
$A_{4,7}$	$f_{14}^1 = 2, f_{24}^2 = 1, f_{34}^2 = 1$ $, f_{34}^3 = 1, f_{23}^1 = 1$	$\begin{pmatrix} a_2^2 & 0 & 0 & 0 \\ -a_2 a_5 & a_2 & 0 & 0 \\ -a_2 a_5 + a_2 a_4 - a_1 a_5 & a_1 & a_2 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,8}$	$f_{24}^2 = 1, f_{34}^3 = -1$ $, f_{23}^1 = 1$	$\begin{pmatrix} a_1 a_2 & 0 & 0 & 0 \\ a_1 a_5 & a_1 & 0 & 0 \\ a_2 a_4 & 0 & a_2 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,9}^b$	$f_{14}^1 = 1 + b, f_{24}^2 = 1$ $, f_{34}^3 = b, f_{23}^1 = 1$	$\begin{pmatrix} a_1 a_2 & 0 & 0 & 0 \\ -a_1 a_5 / b & a_1 & 0 & 0 \\ a_2 a_4 & 0 & a_2 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,9}^1$	$f_{14}^1 = 2, f_{24}^2 = 1$ $, f_{34}^3 = 1, f_{23}^1 = 1$	$\begin{pmatrix} a_1 a_4 - a_2 a_3 & 0 & 0 & 0 \\ a_2 a_6 - a_1 a_7 & a_1 & a_2 & 0 \\ a_4 a_6 - a_3 a_7 & a_3 & a_4 & 0 \\ a_5 & a_6 & a_7 & 1 \end{pmatrix}$
$A_{4,9}^0$	$f_{14}^1 = 1, f_{24}^2 = 1$ $, f_{23}^1 = 1$	$\begin{pmatrix} a_2 a_3 & 0 & 0 & 0 \\ a_1 & a_2 & 0 & 0 \\ a_3 a_5 & 0 & a_3 & 0 \\ a_4 & a_5 & 0 & 1 \end{pmatrix}$
$A_{4,10}$	$f_{24}^3 = -1, f_{34}^2 = 1$ $, f_{23}^1 = 1$	$\begin{pmatrix} a_1^2 + a_2^2 & 0 & 0 & 0 \\ -a_1 a_4 - a_2 a_5 & a_1 & a_2 & 0 \\ a_2 a_4 - a_1 a_5 & -a_2 & a_1 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,11}^a$	$f_{14}^1 = 2a, f_{24}^2 = a, f_{24}^3 = -1$ $, f_{34}^2 = 1, f_{34}^3 = a, f_{23}^1 = 1$	$\begin{pmatrix} a_1^2 + a_2^2 & 0 & 0 & 0 \\ -\frac{a(a_1 a_4) + a(a_2 a_5) + a_2 a_4 - a_1 a_5}{a^2+1} & a_2 & -a_1 & 0 \\ \frac{a(a_2 a_4) - a(a_1 a_5) - a_1 a_4 - a_2 a_5}{a^2+1} & a_1 & a_2 & 0 \\ a_3 & a_4 & a_5 & 1 \end{pmatrix}$
$A_{4,12}$	$f_{14}^2 = -1, f_{13}^1 = 1$ $, f_{24}^1 = 1, f_{23}^2 = 1$	$\begin{pmatrix} a_2 & -a_1 & 0 & 0 \\ a_1 & a_2 & 0 & 0 \\ -a_4 & a_3 & 1 & 0 \\ a_3 & a_4 & 0 & 1 \end{pmatrix}$

$$\Lambda_1 = \mathrm{Rotation}_{xy}\mathrm{Boost}_{xz}\mathrm{Boost}_{yz}C \tag{47}$$

where:

$$\mathrm{Rotation}_{xy} = \begin{pmatrix} \cos(a_1) & \sin(a_1) & 0 & 0 \\ -\sin(a_1) & \cos(a_1) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{48}$$

$$\mathrm{Boost}_{xz} = \begin{pmatrix} \cosh(a_2) & 0 & \sinh(a_2) & 0 \\ 0 & 1 & 0 & 0 \\ \sinh(a_2) & 0 & \cosh(a_2) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{49}$$

$$\mathrm{Boost}_{yz} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cosh(a_3) & \sinh(a_3) & 0 \\ 0 & \sinh(a_3) & \cosh(a_3) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{50}$$

$$C = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & a_4 \end{pmatrix} \tag{51}$$

$$\tag{52}$$

$$\Lambda_2 = \mathrm{Rotation}_{xy}\mathrm{Rotation}_{xz}\mathrm{Rotation}_{yz}C \tag{53}$$

where:

$$\mathrm{Rotation}_{xy} = \begin{pmatrix} \cos(a_1) & \sin(a_1) & 0 & 0 \\ -\sin(a_1) & \cos(a_1) & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{54}$$

$$\mathrm{Rotation}_{xz} = \begin{pmatrix} \cos(a_2) & 0 & -\sin(a_2) & 0 \\ 0 & 1 & 0 & 0 \\ \sin(a_2) & 0 & \cos(a_2) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{55}$$

$$\mathrm{Rotation}_{yz} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos(a_3) & \sin(a_3) & 0 \\ 0 & -\sin(a_3) & \cos(a_3) & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \tag{56}$$

$$C = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & a_4 \end{pmatrix} \tag{57}$$

References

[1] S.J.Gate, C.M.Hull and M.Roček,"*Twisted Multiplets and New Supersymmetric Nonlinear Sigma Model*", Nucl.Phys. B **248**(1984) 157.

[2] S.Lyakhovich and M.Zabzine, "*Poisson geometry of sigma models with extended supersymmetry* ", Phys.Lett. B **548**(2002) 248, [arXiv:hep-th/0210043].

[3] S. Parkhomenko, " *Extended superconformal current algebras and finite-dimensional Manin triples*", Sov. Phys. JETP. 7 (1992),1-3.

[4] U.Lindström and M.Zabzine ,"*D-branes in N=2 WZW models*", arXiv:hep-th/0212042 v2 5 Mar 2003.

[5] H.Samelson," *A class of complex-analytic manifolds*", Port.Math.(1953), 129-132.

[6] A.Morimoto," *Structures complexes sur les groups de Lie semi-simples* ", C.R.**242** (1956), 1101-1103.

[7] J.E.Snow ,"*Invariant Complex Structures on Four Dimensional Solvable Real Lie Groups*", manuscripta math. **66**,(1990) 397-412 .

[8] G.Ovando ,"*Complex,Symplectic and Kähler structures on four dimensional Lie group*",math.DG/0309146 v1,8 sep 2003,
"*Invariant complex structures on solvable real Lie groups*", manuscripta math. **103**,(2000) 19-30 .

[9] J.Patera and P.Winternitz," *Subalgebras of real three- and four dimensional Lie algebras*", J.Math.Phys.**18**(7) (1977), 1449-1456.

[10] T.Christodoulakis, G.O.Papadopoulos and A.Dimakis ,"*Automorphisms of Real 4 Dimensional Lie Algebras and the Invariant Characterization of Homogeneous 4-Spaces*", arXiv:gr-qc/0209042 v2 13 Nov 2002.

[11] A.Andrada, M.L.Barberis, I.G.Dotti, and G.P.Ovando ,"*Product structures on four dimensional solvable Lie algebras*", math/0402234 v1 [math.RA],13 Feb 2004.

[12] L.Hlavaty and L.Snobl," *Classification of Poisson-Lie T-dual models with two-dimensional targets*", arXiv:hep-th/0110139 v2 20 Dec 2001.

[13] A. Rezaei-Aghdam and M. Sephid," *Complex and biHermitian structures on six dimensional real Lie algebras*". work in progress .